NM-AIST Repository

https://dspace.mm-aist.ac.tz

Life sciences and Bio-engineering

Masters Theses and Dissertations [LISBE]

2021-12

Factors associated with hypertension among employees in arusha city, Tanzania

Zubery, Dalahile

NM-AIST

https://doi.org/10.58694/20.500.12479/1501 Provided with love from The Nelson Mandela African Institution of Science and Technology

FACTORS ASSOCIATED WITH HYPERTENSION AMONG EMPLOYEES IN ARUSHA CITY, TANZANIA

Dalahile Zubery

A Dissertation Submitted in Partial Fulfillment of the Requirements for the Degree of Master's in Life Sciences of the Nelson Mandela African Institution of Science and Technology

Arusha, Tanzania

December, 2021

ABSTRACT

Tanzania is one of developing countries with a higher prevalence of hypertension than many other developing countries. Healthcare workers, teachers and bankers are occupation groups exposed to hypertension due to the nature of their work. There is currently limited published data on the burden of hypertension among this group in Tanzania. A descriptive cross-sectional study was therefore conducted and collection of the data was done from August 2019 to February 2020 to identify the predicting factors for hypertension among teachers, bankers and healthcare workers in Arusha city council. A total of 305 working adults aged 18-60 years were involved in the study. A modified World Health Organization (WHO) STEPwise approach to chronic disease risk factor surveillance questionnaire collected information about socio-demographic characteristics, lifestyle behaviors, dietary practices, physical activity level, anthropometric measurements and biochemical measurements. The data collected were calculated and ranked by using WHO guidelines. The overall hypertension prevalence among working adults was 23.8%. Age (Adjusted Odds Ratio [AOR=34.98, 95% CI: 1.30-94.03]), alcohol consumption (AOR=6.55, 95% CI: 1.22-35.28), low salary (AOR=6.44, 95% CI: 1.12-37.18) and high Low-Density Lipoprotein cholesterol (LDL-C) (AOR=5.93, 95% CI: 1.24-28.45), were significantly associated with hypertension. These findings can be used by local policymakers, education, financial and health sectors as baseline information when planning strategies for management and prevention of hypertension and other Non-Communicable Diseases (NCDs) at workplace, by designing workplace wellness programs to mitigate the associated factors.

DECLARATION

I, Dalahile Zubery do hereby declare to the Senate of Nelson Mandela African Institution of Science and Technology that this dissertation is my original work and that it has neither been submitted nor being concurrently submitted for degree award in any other institution.

Dalahile Zubery17/12/2021Candidate NameSignatureDate

The above declaration is confirmed

Dr. Haikael Martin Supervisor (1)

Date

Signature

Prof. Judith Kimiywe

Supervisor (2)

.....

17/12/2021 Date

Signature

COPYRIGHT

This dissertation is copyright material protected under the Berne Convention, the Copyright Act of 1999 and other international and national enactments, on that behalf, on intellectual property. It must not be reproduced by any means, in full or in part, except for short extracts in fair dealing; for researcher private study, critical scholarly review or discourse with an acknowledgement, without written permission of the Deputy Vice-Chancellor for Academic, Research and Innovation, on behalf of both the author and the Nelson Mandela African Institution of Science and Technology.

CERTIFICATION

The undersigned certify that they have read the dissertation titled "Factors Associated with Hypertension among Employees in Arusha City, Tanzania" and recommend for examination in fulfilment of the requirements for the degree of Master's in Life Sciences of the Nelson Mandela African Institution of Science and Technology.

Dr. Haikael Martin

12 202 17

Supervisor 1

Signature

Date

Prof. Judith Kimiywe

N

17/12/2021

Supervisor 2

Signature

Date

ACKNOWLEDGEMENTS

First, I sincerely thanks Allah for strength and sustenance during my study. My appreciation goes to the study respondents for their acceptance to participate in the study, the administrators of banks, healthcare centers and schools where data collection was done. I gratefully acknowledge the laboratory managers of Mount Meru Regional hospital for they are assistance in blood sample analysis.

I sincerely appreciate my supervisors Prof. Judith Kimiywe and Dr. Haikael Martin for their support, advice, guidance, comments, and tireless efforts to assist me during my study. My sincere gratitude goes to CREATES for the financial support which made me accomplish my study.

Finally, my deep gratitude to my parents; my father Mr. Zubery Abdallah Rumisha, my mother Marium Andrew Mapuli, my brothers: Edson Merdard Mapuli and Zavuga Zubery Rumisha for their support and all who directly or indirectly participated in making this research work possible.

DEDICATION

This work is dedicated to my parents.

ABSTR	ACTi
DECLA	RATION ii
COPYR	IGHTiii
CERTIF	FICATION iv
ACKNO	OWLEDGEMENTSv
DEDICA	ATION vi
LIST O	F TABLESx
LIST O	F FIGURES xi
LIST O	F APPENDICES xii
LIST O	F ABBREVIATION AND SYMBOLS xiii
CHAPT	ER ONE1
INTRO	DUCTION1
1.1	Background of the Problem1
1.2	Statement of the Problem2
1.3	Rationale of the Study2
1.4	Objectives
	1.4.1 Main Objective
	1.4.2 Specific Objectives
1.5	Research Questions
1.6	Significance of the Study
1.7	Delineation of the Study4
CHAPT	ER TWO5
LITERA	ATURE REVIEW
2.1	Hypertension or Raised Blood Pressure5
2.2	Prevalence of Hypertension

2.3	Occup	ation and Hypertension	5
2.4	Factors	s Associated with Hypertension	6
	2.4.1	Non-modifiable Risk Factors for Hypertension	6
	2.4.2	Modifiable Risk Factors for Hypertension	6
	2.4.3	The Biochemical Factors Associated with Hypertension	8
СНАРТ	ER THF	REE	9
MATER	RIALS A	ND METHODS	9
3.1	Study 1	Design	9
3.2	Study A	Area	9
3.3	Study 1	Population	9
3.4	Sample	e Size Determination	9
3.5	Sampli	ng Procedure	10
3.6	Data C	ollection Tools	10
3.7	Data C	ollection Methods	10
	3.7.1	Determination of Prevalence of Hypertension	10
	3.7.2	Assessment of Behavioral Factors for Hypertension	11
	3.7.3	The Biochemical Factors Associated with Hypertension	12
3.8	Statisti	cal Analysis	13
3.9	Ethical	Considerations	14
CHAPT	ER FOU	JR	15
RESUL	TS ANE	DISCUSSION	15
4.1	Results	5	15
	4.1.1	Socio-demographic Characteristics of the Study Participants	15
4.2	Prevale	ence of Hypertension	16
4.3	Behavi	oral Risk Factors for Hypertension	17
	4.3.1	Tobacco Use, Alcohol Consumption and Dietary Practices	17

	4.3.2	Physical Activities, Stress Level and Anthropometric Measurement1	7
	4.3.3	Biochemical Factors Associated with Hypertension2	20
4.4	Factors	Associated with Hypertension2	21
4.5	Discuss	sion2	24
СНАРТ	ER FIVI	Ξ2	28
CONCI	LUSION	AND RECOMMENDATIONS2	28
5.1	Conclu	sion2	28
5.2	Recom	mendations2	28
REFER	ENCE		30
APPEN	DICES	4	1
RESEA	RCH OL	JTPUTS	52

LIST OF TABLES

Table 1:	Socio-demographic characteristics of respondents stratified by blood pressure stat	us 16
Table 2:	Prevalence of hypertension	17
Table 3:	Lifestyle factors of respondents stratified by blood pressure status	18
Table 4:	Anthropometric measurement and biochemical factors stratified by blood pressustatus	ire 21
Table 5:	Socio-demographic factors associated with hypertension	22
Table 6:	Lifestyle factors associated with hypertension	23
Table 7:	Anthropometric and biochemical factors associated with hypertension	24

LIST OF FIGURES

Figure 1:	The prevalence	of	underweight,	normal	weight,	overweight	and	obesity	among
	working adults								20

LIST OF APPENDICES

Appendix 1:	Structured questionnaire for the assessment of risk factors for hypertension	on and
	assessment of work-related stress	41
Appendix 2:	Assessment of dietary practices	54
Appendix 3:	Showcard	56
Appendix 4:	Informed consent form	58
Appendix 5:	KNCHREC ethical clearance	61

LIST OF ABBREVIATION AND SYMBOLS

AOR	Adjusted Odds Ratio
BMI	Body Mass Index
BP	Blood Pressure
CI	Confidence Interval
COR	Crude Odds Ratio
CVDs	Cardiovascular diseases
DBP	Diastolic Blood Pressure
HDL-C	High-Density Lipoprotein Cholesterol
JNC VII	Seventh Report of the Joint National Committee on Prevention Detection, Evaluation and Treatment of High Blood Pressure.
KNCHREC	Northern Tanzania Health Research Ethics Committee
LDL-C	Low-Density Lipoprotein cholesterol
MET	Metabolic Equivalent
MmHg	Millimeters of Mercury
Mmol/L	Millimole per Liter
NCDs	Non-Communicable Diseases
OR	Odds Ratio
RBG	Random Blood Glucose
RPM	Revolutions per Minute
S.D	Standard Deviation
SBP	Systolic Blood Pressure
SPSS	Statistical Programme for Social Science
TG	Triglycerides
TC	Total Cholesterol
WC	Waist Circumference
WHO	World Health Organization

CHAPTER ONE

INTRODUCTION

1.1 Background of the Problem

Cardiovascular diseases (CVDs) are the leading cause of death among all non-communicable diseases (NCDs), causing about 44% of all NCDs deaths and 31% of all global deaths (World Health Organization [WHO], 2018). Hypertension is the major risk factor for CVDs, like heart attack, stroke and heart failure (Mozaffarian *et al.*, 2014; Ogah *et al.*, 2012). Globally, 22% of adults have hypertension (WHO, 2018). There is an increase in the number of hypertensive adults from 594 million in 1975 to 1.13 billion in 2015 (Zhou *et al.*, 2017). The highest hypertension prevalence (27%) has been reported in the African region and the lowest prevalence (18%) is found in America (WHO, 2018). In a review of hypertension prevalence in developing countries, it was reported a higher hypertension prevalence in Tanzania than in many other developing countries (Ibrahim & Damasceno, 2012).

Hypertension prevalence in Tanzania has risen from 2% to 10% in 1960s to 13% to 79% in 2016 for rural and urban populations (Isungula & Meda, 2017). The higher hypertension prevalence in Tanzania is attributed to tobacco smoking, excessive use of alcohol, unhealthy eating habits and physical inactivity (Mayige *et al.*, 2011). A sedentary lifestyle is one of predicting factors for hypertension (Henson *et al.*, 2013) ascribed by rapid unplanned urbanization, which influences the adaptation of sedentary occupations (Brownson *et al.*, 2005). Occupation is the predicting factor for hypertension due to its association with socio-economic status and lifestyle factors such as physical inactivity and sedentary lifestyle (Allman Farinelli *et al.*, 2010; Henson *et al.*, 2013).

Healthcare workers, teachers and bankers are occupation groups identified to be exposed to the risk of hypertension (Mohmmedirfan *et al.*, 2011; Shivaramakrishna *et al.*, 2010; Sovova *et al.*, 2014). The nature of their work makes them spend long hours at work doing sedentary activities also the socio-economic status can influence the adaptation of less physical activities. Determining the factors that may predict hypertension among employees in this study was important due to the following reasons: Firstly, protection of available workforce productivity from chronic illnesses; secondly, healthcare workers play a great role in taking care of patients and promoting a healthy lifestyle for patients; thirdly, teachers are role models to students and

may influence the adaption of healthy lifestyle by students and their parents; and lastly, bankers have a great role in building and maintaining the financial sector in the country.

Despite that, there is limited documentation on the predictors of hypertension among adults in Tanzania, and Arusha specifically. Therefore, the current study aimed to identify the factors associated with hypertension among healthcare workers, teachers and bankers in Arusha City Council for designing workplace intervention for management and prevention of hypertension and other NCDs at workplace.

1.2 Statement of the Problem

While the prevalence of hypertension in adults among developed countries has declined in the last few decades, it has been stable or increasing in many developing countries (WHO, 2018). Although there is limited published data on the predictors of hypertension among working adults in Tanzania, the 2012 WHO Stepwise survey reported that 26% of the adult population is hypertensive, with 68% of the population never screened for blood pressure (WHO, 2012). In Tanzania, the prevention and management of NCDs at workplace through workplace programs for physical activity and healthy eating habit is still a challenge.

Workplace health programs have been identified by WHO as the most cost-effective ways for management and prevention of hypertension in countries where the provisions of health care for chronic diseases are still limited (Bosu, 2016; WHO, 2010). To implement workplace health programs, World Health Organization (WHO) recommended identifying the factors associated with hypertension as an important step in providing baseline information in planning for prevention and management of hypertension and other NCDs (WHO, 2013).

Since there is limited information to inform policymakers on the predicting factor for hypertension among working adults, the study aimed to identify the predicting factor for hypertension among healthcare workers, teachers and bankers to generate baseline information in designing workplace intervention for the prevention and management of hypertension and other NCDs at workplace.

1.3 Rationale of the Study

The higher prevalence of hypertension have been reported in Tanzania than in many other developing countries. Despite that, there is limited information on the burden of hypertension

among working adults. Therefore this study aimed to determine the factors associated with hypertension among healthcare workers, teachers and bankers. This will enable the identification of predictors of hypertension among working adults and help to design workplace intervention for the prevention and management of hypertension at workplace.

1.4 Objectives

1.4.1 Main Objective

To determine the factors associated with hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council.

1.4.2 Specific Objectives

- To determine the prevalence of hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council.
- (ii) To assess the behavioral factors associated with hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council.
- (iii) To determine the biochemical factors associated with hypertension such as blood glucose and serum lipid profile among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council.

1.5 Research Questions

- (i) What is the prevalence of hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council?
- (ii) What are the behavioral factors for hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council?
- (iii) What are biochemical factors associated with hypertension among healthcare workers, teachers and bankers aged 18-60 years old in Arusha City Council?

1.6 Significance of the Study

The findings from the current study fill the knowledge gap on the factors associated with hypertension among working adults in Arusha and generate baseline information about the factors associated with hypertension among healthcare workers, teachers and bankers. It informs the policy makers to develop workplace intervention strategies to mitigate the risk factors for hypertension and related NCDs but also indirect community intervention through patients, students and customers.

1.7 Delineation of the Study

The current study was descriptive cross-sectional which is limited on exploring the causal relationship between the risk factors and hypertension since data were collected at once. Furthermore, the study participants were working professionals from urban residents hence, generalizing the result to all working adults is not appropriate especially for those who are working in rural areas. In addition, majority of the study respondents were female in comparison with other studies due to a large proportion of females than males in the study population. Another limitation encountered was on blood sample collection for biochemical assay tests. Random blood samples were collected among study respondents. This may cause misclassification of the biochemical assay, especially on triglyceride levels as it remains high for several hours after a meal. Moreover, the best way to assess dietary intake of salt is by measuring the urinary electrolyte but in this study, the assessment was based on a questionnaire only. Also, the one month recall of foods consumed on assessment of dietary intake may be subjected to recall bias that may result to misreporting or over-reporting.

CHAPTER TWO

LITERATURE REVIEW

2.1 Hypertension or Raised Blood Pressure

Hypertension is a condition in which the blood vessels have persistently raised pressure (WHO, 2013). The measurement of blood pressure is done in millimeters of mercury (mmHg) and recorded as systolic blood pressure, the highest pressure in blood vessels when the heart beats and diastolic blood pressure, the lowest blood pressure in blood vessels between heartbeats (WHO, 2013). The normal blood pressure for adults is systolic blood pressure less than 120 mmHg and diastolic blood pressure less than 80 mmHg and raised blood pressure is categorized as systolic blood pressure equal to or above 140 mmHg and diastolic blood pressure equal to or above 90 mmHg. Normal blood pressure is required for the overall health and wellbeing of vital organs such as heart, kidneys and brain (Momin *et al.*, 2012; WHO, 2013). Hypertension is associated with cardiovascular disease, hardening of the arteries, kidney disease, stroke, and eye damage (WHO, 2009).

2.2 Prevalence of Hypertension

Globally, the hypertension prevalence among adults is 22%, whereby one in five women and one in four men had hypertension (WHO, 2018). In sub-Saharan Africa, the overall prevalence of hypertension ranges from 30.0% to 31.1% among adults indicating a major health threat to the community (Ataklte *et al.*, 2014; Nduka *et al.*, 2015). The hypertension prevalence varies from country to country as follows; Kenya (24.5%), Namibia (46.0%), Ethiopia (35.2%), and Tanzania (25.9%) (Alwan, 2011; Craig *et al.*, 2018; Kagaruki & Mayige, 2016; Mohamed *et al.*, 2018). The variation in hypertension prevalence between countries may be caused by a difference in the level of urbanization, cultural habits, dietary practices and ethnicity (Ogah & Rayner, 2013). Despite, the increased hypertension prevalence in developing countries, including Tanzania, there is limited documentation about the hypertension prevalence among working adults.

2.3 Occupation and Hypertension

Occupation is another factor associated with hypertension (Allman Farinelli *et al.*, 2010; Henson *et al.*, 2013). Healthcare workers, teachers and bankers are recognized as a high-risk

group exposed to hypertension and other cardiovascular diseases (Awosan *et al.*, 2017; Mohmmedirfan *et al.*, 2011; Shivaramakrishna *et al.*, 2010; Sovova *et al.*, 2014). This is caused by sedentary nature of their work that makes them less physically active (Ogungbenle *et al.*, 2017). Besides, urbanization has fueled changes in lifestyle, whereby people have reduced physical activity and changes in dietary habits (Contento, 2010). Although different studies have shown that, healthcare workers (Osei-Yeboah *et al.*, 2018; Sovova *et al.*, 2014), teachers (Fikadu & Lemma, 2016; Wijayathunge & Hettiaratchi, 2017), and bankers (Ismail *et al.*, 2013; Kumar & Sundaram, 2014; Manjula *et al.*, 2016) are high-risk group exposed to hypertension, this occupation group is neglected during research studies in Tanzania.

2.4 Factors Associated with Hypertension

2.4.1 Non-modifiable Risk Factors for Hypertension

Non-modifiable risk factors associated with hypertension include sex, family history, age and genetics (Ibekwe, 2015). Increased age is associated with hypertension for both females and males. Studies have shown that, as the age of the person increases, the chance for increased blood pressure becomes high (Craig *et al.*, 2018; Kunutsor & Powles, 2009; Mohamed *et al.*, 2018). Other studies done in Africa including individuals aged 15 years and above had similar findings (Dzudie *et al.*, 2012; Walker *et al.*, 2000).

High risk of hypertension is mostly found in males compared to females (Asfaw *et al.*, 2018; Ekwunife *et al.*, 2010; Ibrahim *et al.*, 2008) but after menopause, the risk of hypertension in term of age become the same. World Health Organization (WHO) Stepwise survey conducted in African countries between 2003-2009 has found a high hypertension prevalence in males than females in the majority of the countries although in some countries the findings were vice-versa (Van de Vijver *et al.*, 2014). The association between genetics and hypertension depends on the family history of hypertension. He *et al.* (2011) studied genes relating to hypertension and found 20 different genes which affected sodium handling in the kidney hence affecting the blood pressure of the affected individual.

2.4.2 Modifiable Risk Factors for Hypertension

The behavioral factors associated with hypertension include unhealthy eating habits, overweight and obesity, physical inactivity, excessive use of alcohol and tobacco smoking which result in metabolic syndrome such as hyperlipidemia, diabetes and hypertension (WHO,

2013). Unhealthy eating is associated with too much consumption of trans-fats and saturated fat, high dietary intake of salt and low consumption of vegetables and fruits (WHO, 2012). The association between unhealthy eating habit and hypertension have been observed in different studies (Njelekela *et al.*, 2003; Nurwanti *et al.*, 2018; Onyango *et al.*, 2017; Popkin, 2006).

Overweight and obesity are other factors associated with hypertension in the population. Excess body weight increases the demand for nutrients and oxygen to the tissues, the increase in blood volume that flow through blood vessels increase the pressure on the arteries which can result in hypertension (Bouchi *et al.*, 2016). For adults aged 18 years and above the estimate shows that among three adults one of them is overweight and more than one in ten are obese, whereby women have excess body weight compared to men (WHO, 2016).

Physical inactivity is the contributing factor for hypertension. The WHO has reported that 28% of adults have inadequate physical activity according to the recommendation while men are active by 32% compared to females 23% (WHO, 2018). Findings show that physical inactivity contributes to the high risk of hypertension and other heart diseases due to increased heart rate that hardens the contraction of the heart and increases pressure to the arteries also, physical inactivity is associated with excessive weight gain (Hayes *et al.*, 2002; Huai *et al.*, 2013; Ketkar *et al.*, 2015, Werneck *et al.*, 2018).

Another factor associated with hypertension is excessive alcohol consumption which results in disabilities and premature deaths among different populations in the world (WHO, 2008). Excessive alcohol users are more exposed to hypertension compared to non-alcohol users (Kotwani *et al.*, 2013; Mohamed *et al.*, 2018; Wamala *et al.*, 2009; Williams *et al.*, 2013). There is more than one mechanism that links excessive alcohol consumption and hypertension such as baroreceptors sensitivity, the effect on the level of calcium and cortisol. For example, baroreceptors help the body to regulate blood pressure but excessive use of alcohol prevents the detection of the body's baroreceptors to detect the need of stretching the blood vessels to increase the diameter of the blood vessels resulting in raised blood pressure (Husain *et al.*, 2014). The use of tobacco which includes smoking and smokeless tobacco is also associated with onset of hypertension caused by the release of tobacco chemicals that damage the arteries walls and cause narrowing of the arteries which end up with raised blood pressure (Burns, 2003; Stein *et al.*, 2008).

2.4.3 The Biochemical Factors Associated with Hypertension

Diabetes and hyperlipidemia are metabolic syndromes associated with hypertension. Diabetes is characterized by the failure of the pancreas to produce insulin hormone or the pancreas having a reduced ability to produce insulin in the body (Association, 2005). Diabetes and hypertension may influence each other. This is because insulin resistance cause sugar builds up in the blood that increases the volume of the blood and raises the blood pressure. Also, diabetes can damage the small blood vessels resulting in stiffening of blood vessels which results to hypertension (Kabakov *et al.*, 2006). The association between diabetes and hypertension have been reported in different studies (Lloydm Sherlock *et al.*, 2014; Mayige *et al.*, 2011; Mosha *et al.*, 2017).

The raised lipid profile is associated with the formation of atherosclerosis that hardens and narrows the arteries and increases the chance of developing hypertension (Arora *et al.*, 2007; Chamba *et al.*, 2017). Raised lipid profile has been reported in several general population studies in Africa (Vorster, 2002). However, raised lipid profile is increasing in developing countries like Tanzania. A study done by Kagaruki and Mayige (2016) for example, has reported elevated serum Total Cholesterol prevalence of 4.4% and 25.8% prevalence of elevated Triglycerides (TG) among adults in Tanzania. Determining the raised blood glucose and abnormal lipid profile levels among working adults fills the knowledge gap on the relationship between lipid profile and hypertension among working adults.

CHAPTER THREE

MATERIALS AND METHODS

3.1 Study Design

The study was a descriptive cross-sectional that measure the outcome (hypertension) and exposure at one point in time.

3.2 Study Area

The study was conducted in Arusha City Council, Arusha Region. It is a third urbanized region in Tanzania due to an urban population of 31.3% than the national average of urban population 23.1% (Worrall *et al.*, 2017). Arusha City is found near the foot of Mount Meru. As per the National Census of 2012, the city has a total population of 416 442 of which 160 091 are employees (The United Republic of Tanzania [URT], 2016). By comparing with other districts in the region, Arusha city has a high number of employees. The Southern part of the region is bordered by Singinda and Manyara regions, the Northern part is bordered by Kenya, the Eastern part is bordered by Kilimanjaro region and the Western part is Simiyu region. The region is a global tourist destination having national parks such as Lake Manyara National Park, Arusha National park and Ngorongoro Conservation Area (URT, 2015).

3.3 Study Population

The study involved healthcare workers, primary and secondary teachers and bankers aged between 18-60 years working in the Arusha City Council. The study included both males and females from government and non-government institutions excluding part-time workers and pregnant women.

3.4 Sample Size Determination

The sample size was estimated by using the Fischer formula (Fischer et al, 1991).

$$n = \frac{Z^2 p q}{d^2}$$

Where; n = sample size, p = the prevalence of hypertension, d is precision of estimate =5%, Z = standard value of 95% CI =1.96 and q= 1-p.

The minimal sample size estimated was 380 participants; the Prevalence of hypertension was 45%, from a study conducted at Arusha City council by Katalambula *et al.* (2017).

3.5 Sampling Procedure

A multistage sampling technique was used in this study. At the first stage, clusters were formed purposively for Healthcare institutions, Educational institutions (primary and secondary schools) and Financial Institutions (Banks). In the second stage, out of 25 wards, 3 wards (Levolosi, Sekei, and Themi) were selected purposively due to an adequate representative number of health centers, schools and banks. In the third stage, randomly selection of health centers, schools and banks from 3 wards were as follows: out of 21 health centers 8 were selected by using ballot method and lastly, a representative sample was obtained by using probability proportional to the size and the sample from healthcare workers was 51, teachers 176 and bankers 153. From each site, the eligible number of participants from health center was at least 8 healthcare workers, for schools and banks at least 13 participants from each and at this stage the participation was voluntary.

3.6 Data Collection Tools

Participant enrollment and collection of data was done from August 2019 to February 2020. The pre-tested modified WHO STEP-wise survey questionnaire including the Global Physical Activity Questionnaire translated from English to Swahili for easy administration collected data on behavioral factors, which included socio-demographic data, as well as information on alcohol consumption, tobacco use, dietary intake, anthropometric measurement, biochemical measurement, physical activity level and history of chronic diseases (Appendix 1). The dietary habit of the study participants was collected by using the Food Frequency Questionnaire (Appendix 2) which was validated by Jordan *et al.* (2013).

3.7 Data Collection Methods

3.7.1 Determination of Prevalence of Hypertension

Blood pressure was measured by using a digital sphygmomanometer (CITIZEN®) in a seated position. Three measurements were done in the right upper arm after 5 minutes rest interval and the average of it was used in data analysis. The classification of blood pressure based on

the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC VII) as follows: Normotensives were classified as systolic blood pressure (SBP) < 120 mmHg and diastolic blood pressure (DBP) < 80 mmHg, prehypertension (SBP 120–129 mmHg or DBP 80–89 mmHg), hypertension stage 1 (SBP 140–159 mmHg or DBP 90–99 mmHg), and hypertension stage 2 (SBP \geq 160 mmHg or DBP \geq 100 mmHg) (Chobanian *et al.*, 2003). The prevalence of hypertension was determined using the WHO formula (WHO, 2006).

3.7.2 Assessment of Behavioral Factors for Hypertension

(i) Assessment of Tobacco Use and Alcohol Consumption

The use of tobacco was assessed by previous and current smoking status, the duration of smoking, daily frequency of tobacco use, exposure to second-hand smoking and the use of smokeless tobacco. The assessment of alcohol consumption was done by recalling if the study participant was a lifetime abstainer, 12-month abstainer and past 30-day drinker. Also, the frequency of alcohol use, type of alcohol consumed frequently and consumption of alcohol with food was documented.

(ii) Assessment of Dietary Intake

The assessment of dietary intake included the number of days on the consumption of fruit and vegetable in a week, the serving portion, type of meals eaten at work per week, type of fat and oils used frequently in meal preparation at home. Salt consumption assessment was done by asking the study respondents about the frequency of salt used during food preparation at home, the use of table salt, perception of the amount of salt consumed and its association with health problems, also the health benefit of minimizing salt intake and the strategies undertaken to control salt consumption. The Food Frequency Questionnaire (Appendix 2) was used to assess the frequency of consumption of food groups in a month to determine the dietary practices of the participants.

(iii) Anthropometric Measurements

Anthropometric measurements were taken by a qualified healthcare provider. A portable SHORRTM stadiometer (Shorr Productions, Olney, MA, USA), was used to measure the height of the study participants. The respondents removed shoes and stand still with their back against

the wall and heels put together in a V-shape while looking forward and the measurement was taken nearest to 0.5 cm. SECATM digital weighing scale (Seca GmbH, Hamburg, and Germany) was used to measure body weight. The participants were asked to remove their footwear and heavy accessories and measurement were taken nearest 0.1 kg. Then, the Body Mass Index (BMI) was calculated using the WHO formula (kg/m²) and the results were ranked as follows: $\leq 18.4 \text{ kg/m2}$ as underweight, 18.5-24.9 kg/m2 as normal weight, 25.0-29.9 kg/m2 as overweight, and $\geq 30 \text{ kg/m2}$ as obese (WHO, 2016). Waist circumference was measured midway between the last rib bone and iliac bone by using a measuring tape and the measurement was taken nearest to 0.5 cm. The results were ranked as follows: >80 cm for females and >94 cm for males had increased risk of metabolic complication and >88 cm for females and >102 cm for males had substantially increased risk of metabolic complication. The measurements lined with the standard method of anthropometric measurement (WHO, 2005, 2014).

(iv) Assessment of Physical Activity Level and stress

The physical activity level was assessed by using the Global Physical Activity Questionnaire by considering the intensity, frequency and duration of physical activity in different areas specifically during transport, at work and in recreational areas. The collected data included the number of days in a week, time spent doing physical activity in hours and minutes during transport, at work and recreational area for at least 10 minutes or more continuously in a day (Armstrong & Bull, 2006). To determine the Metabolic Equivalent (MET) which measure the energy expenditure of physical activity, the intensity, frequency and duration of physical activities performed in a week were used and physical activity level was classified by following a standard classification as high physical activity level, moderate physical activity level and low physical activity level (Armstrong & Bull, 2006) and sedentary work was classified as a job that needs a person to sit for 6 hours out of 8 hours at work. For stress, the participants were asked about feeling stressed at work, perception of the salary, working for long hours, unrealistic time pressure and respect at work.

3.7.3 The Biochemical Factors Associated with Hypertension

(i) Determination of Blood Glucose and Lipid Profile

A standardized Glucometer machine (GlucoPlusTM Inc. Quebec, Canada) was used to measure Random Blood Glucose (RBG) by using the capillary finger-prick method. Diabetes was determined by having random plasma glucose of $\geq 11.1 \text{ mmol/L}$ or the use of diabetic drugs. A random sub-sample of half of the participants was selected for additional measurement of serum lipid profile. The size of this sub-sample (153 participants) was determined based on the availability of resources. A qualified healthcare provider collected blood samples from participants. The Vacutainer needle collected blood from the median basilic vein of the participant and stored in a red top Vacutainer® blood collection tube and left to clot for 45 minutes. Then, centrifugation was done at a spin rate of 1100-1300 rpm for 10 minutes. Afterwards, serum was pipetted from the clotted blood and stored in a 2 ml Eppendorf tube. All serum samples were frozen at -20 °C until analysis. Serum lipid profiles included; Total cholesterol (TC), Triglyceride (TG), High-Density Lipoprotein Cholesterol (HDL-C) and Low-Density Lipoprotein Cholesterol (LDL-C). All biochemistry assays were carried out at Mount Meru Regional Hospital laboratory using fully automated Chemistry auto analyzer XL-180 (Vital Scientific, B.V. Kanaalweg 24, and Netherlands). The identification of serum lipid profile was ranked according to the WHO cut-off point (WHO, 2005).

3.8 Statistical Analysis

The collected data were entered in Microsoft ExcelTM (2016) and exported to Statistical Package for Social Science Version 20.0 software (SPSS Inc., Chicago, IL, USA) for analysis. Descriptive statistics analyzed social-demographic characteristics and frequency and percentage were used to analyze categorical data. For continuous data mean and standard deviation were used. On the assessment of the difference between groups, a chi-squared test was used for categorical variables and a t-test for continuous variables. To test the normality assumption of variable distribution the Kolmogorov–Smirnov test was used. Binary logistic regression analysis was done to determine the factors that were associated with hypertension. The independent variables were social demographic characteristics, behavioral risk factors which included; dietary habits, dietary salt intake, Body Mass Index, stress, physical activity level, alcohol consumption and tobacco smoking while the dependent variable was hypertension. To test the goodness fit of the model Hosmer-Lemeshow statistic was used and the results reported in odds ratios (OR) at 95% confidence intervals and p < 0.05 was statistically significant.

3.9 Ethical Considerations

The study was conducted according to the Declaration of Helsinki. Ethical clearance certificate number KNCHREC 0014 was obtained from Northern Tanzania Health Research Ethics Committee (KNCHREC). Written permission to conduct research was also sought from Arusha City Council and administrators of the selected banks, schools and health facilities. Respondents signed written informed consent after being briefed about the study. To ensure confidentiality data collection was done individually in a special prepared room in their working area and codes known only by researchers were used instead of names in identifying participants to maintain anonymity.

CHAPTER FOUR

RESULTS AND DISCUSSION

4.1 Results

4.1.1 Socio-demographic Characteristics of the Study Participants

A total of 305 study participants were included in this study. However, 19 were excluded from statistical analysis due to the incompleteness of data. The final sample size was 286 respondents from three occupations: healthcare workers (61; 21.3%), teachers (131; 45.8%), and bankers (94; 32.9%). The majority of the respondents (204; 71.3%) were female. Nearly half of the respondents (121; 42.3%) were aged 30–39 years; 141 (49.3%) of the study participants had attended college education and (182; 63.6%) were married or cohabiting. Nearly half 122 (42.7%) of the study respondents worked for more than seven years in the current institution and in term of estimated monthly income per month, 22 (7.7%) were earning less than 250 000 (Table 1).

Variable	Total	Normotension	Pre-	Hypertension	P-
	n (%)	n (%)	hypertension	n (%)	value
			n (%)		
Profession					
Healthcare workers	61 (21.3)	33 (54.1)	17 (27.9)	11 (18.0)	0.082
Teachers	131 (45.8)	69 (52.7)	37 (28.2)	25 (19.1)	
Bankers	94 (32.9)	42 (44.7)	20 (21.3)	32 (34.0)	
Age					
18-29 Years	75 (26.2)	44 (58.7)	17 (22.7)	14 (18.7)	0.029
30-39 Years	121(42.3)	65 (53.7)	34 (28.1)	22 (18.2)	
40-49 Years	49 (17.1)	19 (38.8)	15 (30.6)	15 (30.66)	
50-60 Years	41 (14.3)	16 (39.0)	8 (19.5)	17 (41.5)	
Gender					
Male	82 (28.7)	37 (45.1)	23 (28.0)	22 (26.8)	0.526
Female	204 (71.3)	107 (52.5)	51 (25.0)	46 (22.5)	
Education level					
College	141 (49.3)	76 (53.9)	37 (26.2)	28 (19.9)	0.637
University graduate	122 (42.7)	57 (46.7)	31 (25.4)	34 (27.9)	
degree	23 (8.0)	11 (47.8)	6 (26.1)	6 (26.1)	
Postgraduate	. ,	× ,		. ,	
degree					
Marital status					
Single	104 (33.2)	57 (60.0)	20 (21.1)	18 (18.9)	0.028
Married/Cohabitati	18 2(63.6)	84 (46.2)	51 (28.0)	47 (25.8)	
ng					
Years with the					
current institution					
Less than a year	24 (8.4)	13 (54.2)	4 (16.7)	7 (29.2)	0.194
1-3 years	58 (20.3)	36 (62.1)	14 (24.1)	8 (13.8)	
4-6 years	82 (28.7)	42 (51.2)	18 (22.0)	22 (26.8)	
7 years and above	122 (42.7)	53 (43.4)	38 (31.1)	31 (25.4)	
The estimated					
income per month					
Less than 250 000	22 (7.7)	12 (54.5)	4 (18.2)	6 (27.3)	0.299
250 000-500 000	79 (27.6)	47 (59.5)	18 (22.8)	14 (17.7)	
500 000-750 000	83 (29.0)	41 (49.4)	25 (30.1)	17 (20.5)	
750 000-1 000 000	64 (22.4)	26 (40.6)	20 (31.3)	18 (28.1)	
More than 1 000	38 (3.3)	18 (47.4)	7 (18.4)	13 (34.2)	
000					

 Table 1: Socio-demographic characteristics of respondents stratified by blood pressure status

4.2 Prevalence of Hypertension

The overall mean (\pm SD) Systolic Blood Pressure (SBP) of the study participants was 124.4 \pm 19.2 mmHg and Diastolic Blood Pressure (DBP) was 79.0 \pm 12.8 mmHg (Table 4). About half (144; 50.3%) of the study participants had normal blood pressure, 24 (8.4%) had

hypertension stage 1 and 14 (4.8%) had hypertension stage 2 (Table 2). For those who had hypertension, 42 (61.8%) were aware of their condition and 25 (59.5%) were using antihypertensive medication. Furthermore, 14 (28.6%) of hypertensive working adults had a family history of hypertension.

Table 2. I revalence of hypertension		
Variable	Ν	Percent (%)
Normal (SBP < 120 and DBP < 80)	144	50.3
Pre-hypertension (SBP 120–139 or DBP 80–89)	74	25.9
Hypertension		
Isolated Systolic hypertension	18	6.3
$(SBP \le 140 \text{ mmHg and } DBP \ge 90 \text{ mmHg})$		
Isolated Diastolic hypertension	12	4.2
(SBP \geq 140 mmHg and DBP < 90 mmHg)		
Stage 1 hypertension (SBP 140–159 or DBP 90–99)	24	8.4
Stage 2 hypertension (SBP \ge 160 or DBP \ge 100)	14	4.9

Table 2: Prevalence of h	ypertension
--------------------------	-------------

4.3 **Behavioral Risk Factors for Hypertension**

4.3.1 **Tobacco Use, Alcohol Consumption and Dietary Practices**

None of the respondents was using tobacco. The overall prevalence of alcohol consumption was 61 (21.3%). More than half of the study participants 180 (62.9%) add table salt during eating. Furthermore, those consuming adequate fruits and vegetables per week were 123 (43.0%) and 144 (50.3%) respectively (Table 3).

4.3.2 **Physical Activities, Stress Level and Anthropometric Measurement**

The majority of the study participants 236 (82.5%) were sedentary workers and a low level of physical activity was prevalent in the study population to about 127 (44.4%). About 115 (40.2%) of the respondents were feeling stressed at work and on the perception of salary meet the daily basic need, majority of them 234 (81.8%) reported that the salary does not meet the daily basic need (Table 3). The overall mean BMI (BMI \pm SD) was 28.4 \pm 5.9 kg/m², and normotension and hypertension participants were 28.7 ± 5 .8 kg/m² and 31.6 ± 5.5 kg/m² respectively. About 89 (31.1%) of the study respondents were overweight and 108 (37.8%) were obese (Fig. 1). The overall mean waist circumference of the study participant was 92.0±12.9 cm and 195 (68.2%) of the participants had abdominal obesity (Table 4).

Variable	Total	Normotension	Pre-hypertension	Hypertension	P-value
variable	n (%)	n (%)	n (%)	n (%)	
Alcohol consumption					
Lifetime abstainers	165 (57.7)	82 (49.7)	51 (30.9)	32 (19.4)	0.009
Past drinkers	60 (21.0)	36 (60.0)	12 (20.0)	12 (20.0)	
Current drinkers	61 (21.3)	26 (42.6)	11 (18.0)	24 (39.3)	
Exposure to second-hand smoke					
No	210 (73.4)	105 (50.0)	58 (27.6)	47 (22.4)	0.450
Yes	76 (26.6)	39 (51.3)	16 (21.1)	21 (27.6)	
Type of oil or fat used at home					
Sunflower oil	204 (71.3)	100 (49.0)	56 (27.5)	48 (23.5)	0.729
Palm oil	60 (21.0)	31 (51.7)	15 (46.7)	14 (23.3)	
No specific cooking oil	22 (7.7)	13 (59.1)	3 (13.6)	6 (27.3)	
Kind of meals eaten at work					
Breakfast	27 (9.4)	13 (48.1)	7 (25.9)	7 (25.9)	0.896
Lunch	114 (39.9)	56 (49.1)	31 (27.2)	27 (23.7)	
Breakfast and lunch	132 (46.2)	69 (52.3)	34 (25.8)	29 (22.0)	
Breakfast, lunch and dinner	13 (4.5)	6 (46.2)	2 (15.4)	5 (38.5)	
Use of table salt	. ,				
Yes	180 (62.9)	98 (54.4)	41 (22.8)	41 (22.8)	0.164
No	106 (37.1)	46 (43.4)	33 (31.1)	27 (25.5)	
Consumption of fruits					
\geq 5 days in a week	123 (43.0)	57 (46.3)	31 (25.2)	35 (28.5)	0.258
< 5 days in a week	163 (57.0)	87 (53.4)	43 (26.4)	33 (20.2)	
Consumption of vegetables					
\geq 5 days in a week	144 (50.3)	75 (52.1)	33 (22.9)	36 (25.0)	0.513
< 5 days in a week	142 (49.7)	69 (48.6)	41 (28.9)	32 (22.5)	
Mode of transport to and from work	× ,	× ,			
Walking	56 (19.6)	33 (58.9)	12 (21.4)	11 (19.6)	0.382
Car and walking	12 (4.2)	4 (33.3)	6 (50.0)	2 (16.7)	
Car and motorcycle	32 (11.2)	15 (46.9)	7 (21.9)	10 (31.3)	
Car	186 (65.0)	92 (49.5)	49 (26.3)	45 (24.2)	

Table 3: Lifestyle factors of respondents stratified by blood pressure status

Variable	Total	Normotension	Pre-hypertension	Hypertension	P-value
	n (%)	n (%)	n (%)	n (%)	
Sedentary work					
Yes	236 (82.5)	118 (50.0)	60 (25.4)	58 (24.6)	0.778
No	50 (17.5)	26 (52.0)	14 (28.0)	10 (20.0)	
Level of physical activity					
High physical activity	34 (11.9)	19 (55.9)	10 (29.4)	5 (14.7)	0.756
Moderate physical activity	125 (43.7)	61 (48.8)	33 (26.4)	31 (24.8)	
Low physical activity	127 (44.4)	64 (50.4)	31 (24.4)	32 (25.2)	
Stress at work					
Yes	115 (40.2)	64 (55.7)	23 (20.0)	28 (24.3)	0.160
No	171 (59.8)	80 (46.8)	51 (29.8)	40 (23.4)	
Working for long hours					
Yes	181 (63.3)	91 (50.3)	43 (23.8)	47 (26.0)	0.397
No	105 (36.7)	53 (50.5)	31 (29.5)	21 (20.0)	
The salary meets the basic need					
Yes	52 (18.2)	23 (44.2)	14 (26.9)	15 (28.8)	0.552
No	234 (81.8)	121 (51.7)	60 (25.6)	53 (22.6)	



Figure 1: The prevalence of underweight, normal weight, overweight and obesity among working adults

4.3.3 Biochemical Factors Associated with Hypertension

The overall mean blood glucose level of the study respondents was 6.1 ± 2.4 (mmol/L). There was a significant difference in the mean level of blood glucose among normotension 5.8 ± 1.8 (mmol/L) and hypertension individuals 7.0 ± 3.4 (mmol/L) (p=0.0001). About 13 (4.5%) of the respondent had raised blood glucose and 11 (84.6%) were using insulin or oral anti-hyperglycemic agent. Those who were aware that they are diabetic were also hypertensive. For serum lipid profile, the mean raised Total Cholesterol (TC) was 5.2 ± 1.5 mmol/L, Triglyceride (TG) 1.8 ± 1.7 mmol/L, raised Low-Density Lipoprotein cholesterol (LDL-C) 2.4 ± 0.8 mmol/L and low High-Density Lipoprotein cholesterol (HDL-C) was 1.2 ± 0.3 mmol/Significantly higher triglyceride (TG) levels were observed among hypertensive compared to normotensive individuals (Table 4).

1				
Variable	Total	Normotension	Hypertension	P-value
Variable	Mean±SD	Mean±SD	Mean±SD	
Height (cm)	161.8 ± 7.4	161.6±7.7	162.2±6.3	0.573
Weight (kg)	76.9±15.4	74.9±15.3	82.9±2.3	0.343
BMI	28.4 ± 5.9	$28.7\pm5~.8$	31.6±5.5	0.530
WC (cm)	29.4 ± 5.8	91.6±11.7	101. 5±11. 4	0.467
SBP (mmHg)	$124.4{\pm}19.2$	116.4±11.6	146.7 ± 19.2	0.024
DBP (mmHg)	79.0±12.8	73.9 ± 8.9	92. 5±10.7	0.356
Blood glucose (mmol/L)	6.1 ± 2.4	5.8 ± 1.8	7.0± 3.4	0.0001
Total cholesterol (mmol/L)	5.2 ± 1.5	5.8 ± 1.8	5.8±1.7	0.060
Triglyceride (mmol/L)	1.8 ± 1.7	$1.7{\pm}1.1$	$2.4{\pm}2.9$	0.0001
Low-Density	2.4 ± 0.8	2.2 ± 0.8	2.7 ± 0.9	0.593
Lipoprotein(LDL-C)				
High-Density	1.2 ± 0.3	1.2±0.3	1.3 ± 0.4	0.538
Lipoprotein(HDL-C)				

 Table 4: Anthropometric measurement and biochemical factors stratified by blood pressure status

4.4 Factors Associated with Hypertension

The results from univariate logistic regression analysis showed age, marital status, alcohol consumption, BMI, abdominal obesity, raised blood glucose, Total Cholesterol (TC) and Low-Density Lipoprotein cholesterol (LDL-C) were predicting factors for hypertension. However, when all the predicting factors were used in multivariable logistic regression analysis, only age, alcohol consumption, low salary and raised Low-Density Lipoprotein cholesterol (LDL-C) were predicting factors for hypertension. In the adjusted odds ratio, the odds of being hypertensive increase as age increases. The odds of working adults aged 30-39 years, 40-49 years and 50-60 years were (AOR=0.86, 95% CI: 0.06-12.57), (AOR=11.07, 95% CI:0.51-24.21) and (AOR=34.98, 95% CI: 1.30-94.03) respectively (Table 5). The results also showed that working adults who consumed alcohol had 6.55 increased odds of being hypertensive (AOR=6.55, 95% CI: 1.22-35.28) compared to non-alcohol users. The odds of study participants who responded that salary does not meet the daily basic needs was 6.44 higher than those who responded salary meets the daily basic need (AOR=6.44, 95% CI: 1.12-37.18 (Table 6). Furthermore, raised Low-Density Lipoprotein cholesterol (LDL-C) was a predicting factor for hypertension. The odds of being hypertensive was 5.93 higher (AOR=5.93, 95% CI: 1.24-28.45) compared to those having normal Low-Density Lipoprotein cholesterol (LDL-C) (Table 7).
Variable	COR (95%CI)	P-value	AOR (95%CI)	P-value
Profession				
Health workers	1		1	
Teachers	1.26 (0.49-3.25)	0.633	1.31 (0.15-11.75)	0.808
Bankers	2.07 (0.76-5.69)	0.156	3.72 (0.26-53.32)	0.334
Age group				
18-29	1		1	
30-39	1.30 (0.41-4.11)	0.660	0.86 (0.06-12.57)	0.912
40-49	5.92 (1.72-20.35)	0.005	11.07 (0.51-24.21)	0.126
50-60	8.27 (2.41-28.42)	0.001	34.98 (1.30-94.03)	0.034
Gender				
Male	1		1	
Female	0.91 (0.39-2.11)	0.827	1.16 (0.13-10.40)	0.894
Education level				
Postgraduate degree	1		1	
University graduate	4.51 (0.27-7.71)	0.662	1.33 (0.07-25.117)	0.851
degree	1.08 (0.21-5.60)	0.930	0.85 (0.04-18.35)	0.918
College				
Marital status				
Single	1		1	
Married/Cohabitating	2.76 (1.17-6.54)	0.021	0.95 (0.14-6.60)	0.959
Years with institution				
Less than a year	1		1	
1-3 years	0.28 (0.40-1.92)	0.195	0.05 (0.002-1.38)	0.076
4-6 years	1.20 (0.29-5.07)	0.800	0.25 (0.01-5.48)	0.382
7 years and above	1.63 (0.41-6.54)	0.493	0.1 (0.004-2.47)	0.161
The estimated income per				
month				
Less than 250 000	1		1	
250 000-500 000	0.71 (0.12-4.12)	0.703	0.31 (0.01-6.42)	0.450
500 000-750 000	1.69 (0.32-8.96)	0.539	4.39 (0.37-51.49)	0.239
750 000-1 000 000	2.63 (0.50-13.92)	0.257	3.53 (0.25-50.32)	0.353
More than 1 000 000	2.25 (0.35-14.61)	0.396	2.24 (0.09-55.43)	0.622

Table 5: Socio-demographic factors associated with hypertension

COR crude odds ratio

AOR adjusted odds ratio

CI confidence intervals

Tuble 0. Enestyle factors assoc	nated with hyper	tension		
Variable	COR (95%CI)	P-value	AOR (95%CI)	P-value
Alcohol consumption				
No	1		1	
Yes	2.59(1.17-5.73)	0.019	6.55(1.22-35.28)	0.029
Exposure to second-hand smoke	1	0.629	1	
No	1.21(0.56-2.64)		2.02(0.35-11.61)	0.433
Yes				
Type of oil or fat used at home				
Sunflower oil	1		1	
Palm oil	0.90 (0.36-2.23)	0.820	2.67 (0.42-16.83)	0.296
No specific oil	1.46 (0.41-5.25)	0.559	1.36 (0.13-14.28)	0.799
Kind of meals eaten at work				
Breakfast	1		1	
Lunch	1.14 (0.33-3.97)	0.834	1.37 (0.19-10.20)	0.757
Breakfast and Lunch	1.40 (0.41-4.83)	0.594	1.55 (0.17-14.47)	0.699
Breakfast, Lunch and Dinner	0.70 (0.06-7.85)	0.772	11.85 (0.23-	0.217
			599.12)	
The use of table salt			,	
No	1		1	
Yes	1.95 (0.88-4.30)	0.098	4.89 (0.87-27.42)	0.071
Consumption of fruits			, ,	
\geq 5 days per week	1		1	
< 5 days per week	0.57 (0.27-1.19)	0.134	0.81 (0.20-3.38)	0.776
Consumption of vegetables	((,	
> 5 days per week	1		1	
< 5 days per week	0.94 (0.45-1.94)	0.860	0.81 (0.18-3.68)	0.789
Mode of transport to and from	(, , , , , , , , , , , , , , , , , , ,		(,	
work	1		1	
Walking	0.60 (0.06-5.93)	0.662	0.55 (0.01-34.82)	0.776
Car and walking	1.50 (0.45-5.03)	0.511	1.13 (0.10-12.61)	0.924
Car and motorcycle	0.99 (0.39-2.50)	0.976	0.30 (0.05-1.77)	0.182
Car	(,	
Sedentary work				
No	1		1	
Yes	1.23 (0.49-3.14)	0.661	0.76 (0.10-3.25)	0.556
Level of physical activity				
High activity level	1		1	
Moderate activity level	1.69 (0.21-2.28)	0.545	0.76 (0.09-6.79)	0.804
Low activity level	0.93 (0.29-3.00)	0.904	0.37 (0.04-3.21)	0.364
Stress at work	0.50 (0.25 0.000)	0.701		0.001
No	1		1	
Yes	0 85 (0 40-1 77)	0.656	0.63 (0.15-2.66)	0 524
Working for long hours	0.00 (0.10 1.77)	0.020	0.00 (0.10 2.00)	0.021
No	1		1	
Yes	0.98(0.47-2.08)	0 964	0 79 (0 06-9 11)	0 747
The salary meets the basic need	5.76 (0.77 2.00)	0.20 f	0.77 (0.00 7.11)	5.7 17
Ves	1		1	
No	128(051-321)	0.603	6 44 (1 12-37 18)	0.037
Family history of hypertension	1.20 (0.31-3.21)	0.005	0.12^{-3}	0.037
No	1		1	
Ves	1 07 (0 40 2 32)	0.865	1 0 42 (0 12 1 52)	0 186
	1.07 (0.49-2.32)	0.005	0.42(0.12-1.32)	0.100

Table 6: Lifestyle factors associated with hypertension

COR crude odds ratio

AOR adjusted odds ratio

CI confidence intervals

Variable	COR (95%CI)	P-value	AOR (95%CI)	P-value
Total cholesterol (TC)				
Normal	1		1	
Raised	2.35 (1.10-5.03)	0.028	2.92 (0.49-17.39)	0.239
Low-Density Lipoprotein(LDL-C)				
Normal	1		1	
Raised	3.40 (1.56-7.40)	0.002	5.93 (1.24-28.45)	0.026
High-Density Lipoprotein(HDL-C)				
Normal	1		1	
Low	0.593 (0.28-1.25)	0.168	0.45 (0.12-1.77)	0.256
Triglyceride (TG)				
Normal	1		1	
Raised	1.56 (0.75-3.27)	0.235	0.63 (0.11-3.60)	0.604
Blood glucose (mmol/L)				
Normal	1		1	
Raised	5.39 (1.23-23.74)	0.026	3.63 (0.11-3.60)	0.366
Body Mass Index				
Normal	1		1	
Overweight	3.25 (0.85-12.43)	0.085	0.71 (0.06-9.11)	0.792
Obesity	5.35 (1.47-19.44)	0.011	2.64 (0.25-28.28)	0.422
Adnominal obesity				
Normal	1		1	
Abdominal obesity	3.46 (1.14-10.52)	0.029	2.19 (0.13-37.48)	0.589
COR crude odds ratio				

Table 7: Anthropometric and biochemical factors associated with hypertension

AOR adjusted odds ratio

CI confidence intervals

4.5 Discussion

The study aimed to identify the predicting factors for hypertension among healthcare workers, teachers and bankers in Arusha city council. Results from this study may have important inferences on strategies for management and prevention of hypertension and other NCDs at workplace. The overall hypertension prevalence among working adults in Arusha City Council was 23.8%, consistent with the Tanzanian national prevalence of hypertension 26% (Kagaruki & Mayige, 2016). Similar findings have been reported in several general population studies conducted in Tanzania 28.0%, Ethiopia 21.0%, Kenya 24.5% and Ghana 28.1% (Dosoo *et al.*, 2019; Fikadu & Lemma, 2016; Galson *et al.*, 2017; Mohamed *et al.*, 2018). However, the prevalence was lower compared with several general population studies done in Tanzania 45%, Zimbabwe 38.4% Turkey 44.0% and South Africa 52% (Erem *et al.*, 2009; Katalambula *et al.*, 2017; Marwiro, 2012; Monakali *et al.*, 2018). The disparities in hypertension prevalence among studies are due to the features of the study population. The current study included healthcare workers, teachers and bankers only while most of the comparative studies were general population studies and there are limited comparable studies of factors associated with

hypertension among working adults. For example, the study done by Katalambula *et al.* (2017) was a population-based descriptive study with a limited number of working adults while the one in Zimbabwe was based on employees working in the city council without specifying the selected departments (Marwiro, 2012).

The current study revealed, an insignificant association between occupation and hypertension. However, there are limited studies to compare three population groups on the association between occupation and hypertension. A study done in Ethiopia, among teachers and bankers reported similar findings in the comparison with two population groups (Fikadu & Lemma, 2016). The current study identified a high prevalence of hypertension among bankers 47.1% compared to teachers 36.8% and healthcare workers 16.2%. This finding was different from a study done in Ethiopia, which reported a high hypertension prevalence among teachers 21.8% than bankers 19.13% (Fikadu & Lemma, 2016). However, the author did not mention the reason for higher hypertension prevalence among teachers than bankers. The higher prevalence of hypertension among bankers in the current study could be caused by sedentary nature of their work as they spend most of the time sitting compared to other occupations and they also have low physical activities that can influence excessive weight gain.

On the other hand, age was a predicting factor for hypertension. The odds of hypertension increased as age increased. Similar findings have been reported in Ghana, India, Nigeria and Bangladesh (Addo *et al.*, 2008; Barua *et al.*, 2018; Kishore *et al.*, 2016; Ulasi *et al.*, 2010). Besides, Dosoo *et al.* (2019) identified an increase in hypertension prevalence with age and the participants who were aged ≥ 60 had higher odds of being hypertensive than other age groups. It has been evident that age is the main non-modifiable risk factor associated with hypertension ascribed by stiffening of the arteries due to the results of the ageing process which predispose the elderly to the risk of developing hypertension (Ferreira *et al.*, 2012; Sun, 2015).

Furthermore, alcohol consumption was a predicting factor for hypertension. These findings are comparable with the studies done in Northern Tanzania, Rural India, urban Varanasi India, Ethiopia and Kenya they reported that hypertension was significantly associated with excessive alcohol consumption (Galson *et al.*, 2017; Kiber *et al.*, 2019; Kishore *et al.*, 2016; Mohamed *et al.*, 2018; Singh *et al.*, 2017). In addition, the result of this study is consistent with two different studies done in Kilimanjaro, Tanzania by Galson *et al.* (2017) and Mitsunaga *et al.* (2008) who reported that alcohol consumption was an important risk factor for hypertension in northern regions of Tanzania due to cultural practices of home-brewing. Cultural practices

which allow the use of local traditional alcohol drinks could be the reason for excessive alcohol consumption in the study population and influenced the development of hypertension as explained by Husain *et al.* (2014). Hence Cultural practices should be considered when implementing a workplace wellness program at workplace especially in the northern regions of Tanzania.

On the assessment of stress as a predicting factor for hypertension, the working adults who reported that salary does not meet the daily basic needs were more likely to be hypertensive than those who reported salary meets the daily basic needs. These results concur with studies done in Japan, Bangladesh and the United State to assess household income and it is association with hypertension among employees. They reported that household income is associated with development of hypertension due to financial stress for fulfilling their basic needs (Barua et al., 2018; Leigh & Du, 2012; Yanagiya et al., 2020). Furthermore, Leigh and Du (2012) found that doubling the wages decrease the risk of hypertension from 30% to 25% for young working adults and 35% to 30% for women. In addition, a study done in Arusha region among pastoralists in Monduli District reported that having no income was associated with hypertension in this pastoralist community due to financial stress to meet the daily basic need (Khamis et al., 2020). However, the current findings were different from other studies done in Ethiopia, China and South Africa which reported that high income is linked with the development of hypertension especially in developing countries by influencing the adaptation of unhealthy lifestyle such as the purchase of calorie-dense foods and adaptation of sedentary lifestyle as underlying causes of hypertension, however, in developed countries high income is protective due to adaptation of healthy lifestyle (Cois & Ehrlich, 2014; Fikadu & Lemma, 2016; Yu et al., 2000). Inadequate salary to meet the daily basic needs among the study population might be the reason for financial stress for the fulfilment of daily basic needs and expose these working adults to the development of hypertension.

Nonetheless, the findings from this study showed that study subjects who had raised Low-Density Lipoprotein cholesterol (LDL-C) had increased odds of being hypertensive than those who had normal Low-Density Lipoprotein cholesterol (LDL-C). This result concurs with that of Akpa *et al.* (2006) and Osuji *et al.* (2012) that, Low-Density Lipoprotein (LDL) was higher among those who had hypertension than normotensive study participants. The odds of being hypertensive in this study was higher than the study done in Bangladesh and Mongolia (Bayart *et al.*, 2018; Choudhury *et al.*, 2014). The raised Low-Density Lipoprotein Cholesterol (LDL-C) among the study participants might be attributed to sedentary nature of their work that makes them less physically active and it is known that doing physical activity can lower the level of Low-Density Lipoprotein Cholesterol (Szapary *et al.*, 2003). Another contributing factor on raised Low-Density Lipoprotein in this study population could be the dietary habit since majority of them reported to consume fried foods at workplace however the consumption of fried food was not significantly associated with hypertension.

The metabolic syndrome such as raised blood sugar, raised cholesterol, abdominal obesity, overweight and obesity were assessed in the current study however only raised Low-Density Lipoprotein was significantly associated with hypertension. Raised blood glucose was associated with hypertension in bivariate analysis but statistically insignificant in multivariate analysis. Also, BMI and abdominal obesity were insignificantly associated with hypertension beside been evidenced in several studies as factors associated with hypertension (Bayray *et al.*, 2018; Gebrihet *et al.*, 2017; Kayima *et al.*, 2015).

CHAPTER FIVE

CONCLUSION AND RECOMMENDATIONS

5.1 Conclusion

In the current study age, alcohol consumption, low salary and raised Low-Density Lipoprotein cholesterol (LDL-C) were the factors associated with hypertension. The prevalence of hypertension, overweight and obesity was high among working adults and the higher prevalence was among bankers compared with other occupations. The finding shows an emerging problem in the working population, especially among bankers as they are exposed to sedentary activities compared to other occupations. Hence this findings inform the policymakers, education, financial and health sectors to design workplace wellness programs that focus on the provision of education about the adaptation of healthy lifestyles especially on physical activities and healthy eating habits to reduce the exposure to risk factors for hypertension and other NCDs at their workplace.

5.2 **Recommendations**

- (i) The policymakers, health sector, financial institutions and education sector should use these findings as baseline information in planning strategies for management and prevention of hypertension and other NCDs at workplace by designing workplace wellness programs like providing education about the adaptation of healthy eating habits and ensuring the accessibility of healthy foods at workplace, weight management programs and stress management program.
- (ii) Further research is recommended on the assessment of salt intake by measuring urinary electrolytes to get the accurate dietary intake of salt instead of depending on the questionnaire only since the majority of respondents were using table salt, and excess salt intake is associated with hypertension.
- (iii) Further research is recommended on the assessment of dietary practices and their association with hypertension since the current study did not find the association between the dietary practices and hypertension as the current study did not collect the quantity of foods consumed. The further linkage between hypertension, other studied

modifiable factors and NCDs should be assessed in the study area as possible causal factors for NCDs.

REFERENCE

- Addo, J., Smeeth, L., & Leon, D. A. (2008). Prevalence, detection, management, and control of hypertension in Ghanaian civil servants. *Ethnicity and Disease*, *18*(4), 505-511.
- Akpa, M., Agomouh, D., & Alasia, D. (2006). Lipid profile of healthy adult Nigerians in Port Harcourt, Nigeria. Nigeria Journal of Medicine, 15(2), 137-140.
- Allman Farinelli, M. A., Chey, T., Merom, D., & Bauman, A. E. (2010). Occupational risk of overweight and obesity: An analysis of the Australian Health Survey. *Journal of Occupational Medicine and Toxicology*, 5(1), 1-9.
- Alwan, A. (2011). *Global status report on non-communicable diseases*. World Health Organization. https://www.who.int/nmh/publications/ncd_report_full_en.pdf.
- Armstrong, T., & Bull, F. (2006). Development of the World Health Organization Global Physical Activity Questionnaire. *Journal of Public Health*, *14*(2), 66-70.
- Arora, M., Koley, S., Gupta, S., & Sandhu, J. (2007). A study on lipid profile and body fat in patients with diabetes mellitus. *The Anthropologist*, 9(4), 295-298.
- Asfaw, L. S., Ayanto, S. Y., & Gurmamo, F. L. (2018). Hypertension and its associated factors in Hosanna town, Southern Ethiopia: Community based cross-sectional study. *BMC Research Notes*, 11(1), 1-6.
- Association, A. D. (2005). Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 28(S37), S5-S10.
- Ataklte, F., Erqou, S., Kaptoge, S., Taye, B., Echouffo Tcheugui, J., & Kengne, A. (2014). Burden of undiagnosed hypertension in sub-Saharan Africa: A systematic review and meta-analysis. *Hypertension*, 65(2), 291-298.
- Awosan, K. J., Adeniyi, S. A., Bello, H., & Bello-Ibrahim, Z. (2017). Nutritional status, weight perception and weight control practices among office employees in Sokoto, Nigeria. *The Pan African Medical Journal*, 27, 279-290.

- Barua, R., Alam, M., Parvin, N., & Chowdhury, R. (2018). Prevalence of hypertension and its risk factors among school teachers in Dhaka, Bangladesh. *International Journal of Research Medical Sciences*, 6(9), 2902-2910.
- Bayart, N. E., Batsaikhan, S., Yondonjamts, P., Dansran, P., Lkhaasuren, G., Turbat, B., Tserendorj, B., & Batjargal, S. (2018). Serum Lipid Profile and Its Association with Hypertension among Mongolian Adults. *Central Asian Journal of Medical Sciences*, 4(3), 203-210.
- Bayray, A., Meles, K. G., & Sibhatu, Y. (2018). Magnitude and risk factors for hypertension among public servants in Tigray, Ethiopia: A cross-sectional study. *PLoS One*, 13(10), 1-10.
- Bosu, W. K. (2016). Determinants of mean blood pressure and hypertension among workers in west africa. *International Journal of Hypertension*, 2016, 1-20.
- Bouchi, R., Ohara, N., Asakawa, M., Nakano, Y., Takeuchi, T., Murakami, M., Sasahara, Y., Numasawa, M., Minami, I., & Izumiyama, H. (2016). Is visceral adiposity a modifier for the impact of blood pressure on arterial stiffness and albuminuria in patients with type 2 diabetes? *Cardiovascular Diabetology*, 15(1), 1-10.
- Brownson, R. C., Boehmer, T. K., & Luke, D. A. (2005). Declining rates of physical activity in the United States: What are the contributors? *Annual Revision Public Health*, 26, 421-443.
- Chamba, N. G., Shao, E. R., Sonda, T., & Lyarru, I. A. (2017). Lipid profile of type 2 diabetic patients at a tertiary hospital in Tanzania: Cross sectional study. *Journal of Endocrinology Diabetes*, 4(1), 1-6.
- Chobanian, A. V., Bakris, G. L., Black, H. R., Cushman, W. C., Green, L. A., Izzo Jr, J. L., Jones, D. W., Materson, B. J., Oparil, S., & Wright Jr, J. T. (2003). The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Jama*, 289(19), 2560-2571.
- Choudhury, K. N., Mainuddin, A., Wahiduzzaman, M., & Islam, S. M. S. (2014). Serum lipid profile and its association with hypertension in Bangladesh. *Vascular Health and Risk Management*, 10, 327-333.

- Cois, A., & Ehrlich, R. (2014). Analysing the socio-economic determinants of hypertension in South Africa: A structural equation modelling approach. *BMC Public Health*, 14(1), 1-11.
- Contento, I. R. (2010). *Nutrition education linking research, theory, and practice*. Jones & Bartlett Publishers.
- Craig, L. S., Gage, A. J., & Thomas, A. M. (2018). Prevalence and predictors of hypertension in Namibia: A national-level cross-sectional study. *PLoS One*, 13(9), 1-14.
- Dosoo, D. K., Nyame, S., Enuameh, Y., Ayetey, H., Danwonno, H., Twumasi, M., Tabiri, C., Gyaase, S., Lip, G. Y., & Owusu Agyei, S. (2019). Prevalence of hypertension in the middle belt of Ghana: A community-based screening study. *International Journal of Hypertension*, 2019, 1-8.
- Dzudie, A., Kengne, A. P., Muna, W. F., Ba, H., Menanga, A., Kouam, C. K., Abah, J., Monkam, Y., Biholong, C., & Mintom, P. (2012). Prevalence, awareness, treatment and control of hypertension in a self-selected sub-Saharan African urban population: A cross-sectional study. *BMJ Open*, 2(4), 1-10.
- Ekwunife, O. I., Udeogaranya, P. O., & Nwatu, I. L. (2010). Prevalence, awareness, treatment and control of hypertension in a nigerian population. *Health*, 2(7), 731-735.
- Erem, C., Hacihasanoglu, A., Kocak, M., Deger, O., & Topbas, M. (2009). Prevalence of prehypertension and hypertension and associated risk factors among Turkish adults: Trabzon Hypertension Study. *Journal of Public Health*, 31(1), 47-58.
- Ferreira, I., Van De Laar, R. J., Prins, M. H., Twisk, J. W., & Stehouwer, C. D. (2012). Carotid stiffness in young adults: A life-course analysis of its early determinants: The Amsterdam Growth and Health Longitudinal Study. *Hypertension*, 59(1), 54-61.
- Fikadu, G., & Lemma, S. (2016). Socio-economic status and hypertension among teachers and bankers in Addis Ababa, Ethiopia. *International Journal of Hypertension*, 2016, 1-8.
- Galson, S. W., Staton, C. A., Karia, F., Kilonzo, K., Lunyera, J., Patel, U. D., Hertz, J. T., & Stanifer, J. W. (2017). Epidemiology of hypertension in Northern Tanzania: A community-based mixed-methods study. *BMJ Open*, 7(11), 1-10.

- Gebrihet, T. A., Mesgna, K. H., Gebregiorgis, Y. S., Kahsay, A. B., Weldehaweria, N. B., & Weldu, M. G. (2017). Awareness, treatment, and control of hypertension is low among adults in Aksum town, Northern Ethiopia: A sequential quantitative-qualitative study. *PLoS One*, *12*(5), 1-16.
- Hayes, L., White, M., Unwin, N., Raj, B., Fischbacher, C., Harland, J., & Alberti, K. (2002). Patterns of physical activity and relationship with risk markers for cardiovascular disease and diabetes in Indian, Pakistani, Bangladeshi and European adults in a UK population. *Journal of Public Health Medicine*, 24(3), 170-178.
- He, F. J., Burnier, M., & MacGregor, G. A. (2011). Nutrition in cardiovascular disease: Salt in hypertension and heart failure. *European Heart Journal*, *32*(24), 3073-3080.
- Henson, J., Yates, T., Biddle, S. J., Edwardson, C. L., Khunti, K., Wilmot, E. G., Gray, L. J., Gorely, T., Nimmo, M. A., & Davies, M. J. (2013). Associations of objectively measured sedentary behaviour and physical activity with markers of cardiometabolic health. *Diabetologia*, 56(5), 1012-1020.
- Huai, P., Xun, H., Reilly, K. H., Wang, Y., Ma, W., & Xi, B. (2013). Physical activity and risk of hypertension: A meta-analysis of prospective cohort studies. *Hypertension*, 62(6), 1021-1026.
- Husain, K., Ansari, R. A., & Ferder, L. (2014). Alcohol-induced hypertension: Mechanism and prevention. *World Journal of Cardiology*, *6*(5), 245-252.
- Ibekwe, R. (2015). Modifiable risk factors of hypertension and socio demographic profile in Oghara, Delta state: Prevalence and correlates. *Annals of Medical and Health Sciences Research*, 5(1), 71-77.
- Ibrahim, M. M., & Damasceno, A. (2012). Hypertension in developing countries. *The Lancet*, *380*(9841), 611-619.
- Ibrahim, N. K., Hijazi, N. A., & AlBar, A. A. (2008). Prevalence and determinants of prehypertension and hypertension among preparatory and secondary school teachers in Jeddah. *Journal of Egypt Public Health Association*, 83(14), 183-203.

- Ismail, I. M., Kulkarni, A. G., Kamble, S. V., Borker, S. A., Rekha, R., & Amruth, M. (2013). Prevalence of hypertension and its risk factors among bank employees of Sullia Taluk, Karnataka. Sahel Medical Journal, 16(4), 139-143.
- Isungula, K., & Meda, J. (2017). The Burden of Hypertension in the Rural and Urban populations of Tanzania: A review of Trends, Impacts and Response. *Tanzania Journal of Health Sciences*, 1(1), 41-52.
- Jordan, I., Hebestreit, A., Swai, B., & Krawinkel, M. B. (2013). Dietary patterns and breast cancer risk among women in northern Tanzania: A case–control study. *European Journal of Nutrition*, 52(3), 905-915.
- Kabakov, E., Norymberg, C., Osher, E., Koffler, M., Tordjman, K., Greenman, Y., & Stern, N. (2006). Prevalence of hypertension in type 2 diabetes mellitus: Impact of the tightening definition of high blood pressure and association with confounding risk factors. *Journal of the Cardiometabolic Syndrome*, 1(2), 95-101.
- Kagaruki, G., & Mayige, M. (2016). Tanzania Steps Survey Report: Ministry of Health and Social Welfare and National Institute for Medical Research. *ResearchGate*, 2016, 1-154.
- Katalambula, L., Meyer, D., Ngoma, T., Buza, J., Mpolya, E., Mtumwa, A., & Petrucka, P. (2017). Dietary pattern and other lifestyle factors as potential contributors to hypertension prevalence in Arusha City, Tanzania: A population-based descriptive study. *BMC Public Health*, 17(1), 1-7.
- Kayima, J., Nankabirwa, J., Sinabulya, I., Nakibuuka, J., Zhu, X., Rahman, M., Longenecker, C. T., Katamba, A., Mayanja Kizza, H., & Kamya, M. R. (2015). Determinants of hypertension in a young adult Ugandan population in epidemiological transition the MEPI-CVD survey. *BMC Public Health*, 15(1), 1-9.
- Ketkar, A. R., Veluswamy, S. K., Prabhu, N., & Maiya, A. G. (2015). Screening for noncommunicable disease risk factors at a workplace in India: A physiotherapy initiative in a healthcare setting. *Hong Kong Physiotherapy Journal*, 33(1), 3-9.

- Khamis, A. G., Senkoro, M., Mwanri, A. W., Kreppel, K., Mfinanga, S. G., Bonfoh, B., & Kwesigabo, G. (2020). Prevalence and determinants of hypertension among pastoralists in Monduli District, Arusha region in Tanzania: A cross-sectional study. *Archives of Public Health*, 78(1), 1-12.
- Kiber, M., Wube, M., Temesgen, H., Woyraw, W., & Belay, Y. A. (2019). Prevalence of hypertension and its associated factors among adults in Debre Markos Town, Northwest Ethiopia: Community based cross-sectional study. *BMC Research Notes*, 12(1), 1-6.
- Kishore, J., Gupta, N., Kohli, C., & Kumar, N. (2016). Prevalence of hypertension and determination of its risk factors in rural Delhi. *International Journal of Hypertension*, 2016, 1-7.
- Kotwani, P., Kwarisiima, D., Clark, T. D., Kabami, J., Geng, E. H., Jain, V., Chamie, G., Petersen, M. L., Thirumurthy, H., & Kamya, M. R. (2013). Epidemiology and awareness of hypertension in a rural Ugandan community: A cross-sectional study. *BMC Public Health*, 13(1), 1151-1161.
- Kumar, S. G., & Sundaram, N. D. (2014). Prevalence and risk factors of hypertension among bank employees in urban Puducherry, India. *International Journal of Occupation*, *Environment Medicine*, 5(2), 94-100.
- Kunutsor, S., & Powles, J. (2009). Descriptive epidemiology of blood pressure in a rural adult population in Northern Ghana. *Rural and Remote Health*, *9*(2), 1095-1107.
- Leigh, J. P., & Du, J. (2012). Are low wages risk factors for hypertension? *The European Journal of Public Health*, 22(6), 854-859.
- Lloyd Sherlock, P., Beard, J., Minicuci, N., Ebrahim, S., & Chatterji, S. (2014). Hypertension among older adults in low-and middle-income countries: Prevalence, awareness and control. *International Journal of Epidemiology*, 43(1), 116-128.
- Manjula, D., Sahu, B., Sasikumar, N., & Babu, G. (2016). Prevalence of Hypertension in school teachers of Bengaluru. *Hypertension*, 1(2), 1-7.

- Marwiro, A. (2012). Prevalence and risk factors for hypertension among Bulawayo City Council employees [Masters Thesis, University of Zimbabwe]. http://citeseerx.ist. psu.edu/viewdoc/download?doi=10.1.1.916.7255&rep=rep1&type=pdf.
- Mayige, M., Kagaruki, G., Ramaiya, K., & Swai, A. (2011). Non-communicable diseases in Tanzania: A call for urgent action. *Tanzania Journal of Health Research*, 13(5), 378-386.
- Mitsunaga, T., & Larsen, U. (2008). Prevalence of and risk factors associated with alcohol abuse in Moshi, Northern Tanzania. *Journal of Bio-social Science*, 40(3), 379-399.
- Mohamed, S. F., Mutua, M. K., Wamai, R., Wekesah, F., Haregu, T., Juma, P., Nyanjau, L., Kyobutungi, C., & Ogola, E. (2018). Prevalence, awareness, treatment and control of hypertension and their determinants: Results from a national survey in Kenya. *BMC Public Health*, 18(3), 1-10.
- Mohmmedirfan, M., Desai, V., & Kavishwar, A. (2011). A study on effect of life style risk factors on prevalence of hypertension among white collar job people of Surat. *The Internet Journal of Occupational Health*, *1*(3), 1-10.
- Momin, M. H., Desai, V. K., & Kavishwar, A. B. (2012). Study of socio-demographic factors affecting prevalence of hypertension among bank employees of Surat City. *Indian Journal of Public Health*, 56(1), 44-48.
- Monakali, S., Goon, D. T., Seekoe, E., & Owolabi, E. O. (2018). Prevalence, awareness, control and determinants of hypertension among primary health care professional nurses in Eastern Cape, South Africa. *African Journal of Primary Health Care and Family Medicine*, 10(1), 1-5.
- Mosha, N. R., Mahande, M., Juma, A., Mboya, I., Peck, R., Urassa, M., Michael, D., & Todd, J. (2017). Prevalence, awareness and factors associated with hypertension in North West Tanzania. *Global Health Action*, 10(1), 1-10.
- Mozaffarian, D., Fahimi, S., Singh, G. M., Micha, R., Khatibzadeh, S., Engell, R. E., Lim, S., Danaei, G., Ezzati, M., & Powles, J. (2014). Global sodium consumption and death from cardiovascular causes. *New England Journal of Medicine*, 371(7), 624-634.

- Nduka, C., Sarki, A., Uthman, O., & Stranges, S. (2015). Impact of antiretroviral therapy on serum lipoprotein levels and dyslipidemias: A systematic review and meta-analysis. *International Journal of Cardiology*, 199, 307-318.
- Njelekela, Sato, T., Nara, Y., Miki, T., Kuga, S., Noguchi, T., Kanda, T., Yamori, M., Ntogwisangu, J., & Masesa, Z. (2003). Nutritional variation and cardiovascular risk factors in Tanzania rural-urban difference. *South African Medical Journal*, 93(4), 295-299.
- Nurwanti, E., Uddin, M., Chang, J. S., Hadi, H., Syed Abdul, S., Su, E., Nursetyo, A., Masud, J., & Bai, C. H. (2018). Roles of sedentary behaviors and unhealthy foods in increasing the obesity risk in adult men and women: A cross-sectional national study. *Nutrients*, 10(6), 704-716.
- Ogah, O. S., Okpechi, I., Chukwuonye, I. I., Akinyemi, J. O., Onwubere, B. J., Falase, A. O., Stewart, S., & Sliwa, K. (2012). Blood pressure, prevalence of hypertension and hypertension related complications in Nigerian Africans: A review. *World Journal of Cardiology*, 4(12), 327-340.
- Ogah, O. S., & Rayner, B. L. (2013). Recent advances in hypertension in sub-Saharan Africa. *Heart*, 99(19), 1390-1397.
- Ogungbenle, E. R., Omeonu, P. E., Aduroja, P. E., & Kukoyi, O. B. (2017). Workplace Health Promotion in Selected Banks of Mushin Local Government Area, Nigeria. *Journal of Scientific Research and Reports, 16*(6), 1-8.
- Onyango, M. J., Kombe, I., Nyamongo, D. S., & Mwangi, M. (2017). A study to determine the prevalence and factors associated with hypertension among employees working at a call centre Nairobi Kenya. *The Pan African Medical Journal*, *5*(27), 178-186.
- Osei Yeboah, J., Kye-Amoah, K. K., Owiredu, W. K., Lokpo, S. Y., Esson, J., Bella-Johnson, B., Amoah, P., & Asumbasiya-Aduko, R. (2018). Cardiometabolic risk factors among healthcare workers: A cross-sectional study at the Sefwi-Wiawso Municipal Hospital, Ghana. *Bio-Medical Research International*, 2018, 1-10.

- Osuji, C. U., Omejua, E. G., Onwubuya, E. I., & Ahaneku, G. I. (2012). Serum lipid profile of newly diagnosed hypertensive patients in Nnewi, South-East Nigeria. *International Journal of Hypertension*, 2012, 1-7.
- Popkin, B. M. (2006). Global nutrition dynamics: The world is shifting rapidly toward a diet linked with non-communicable diseases. *The American Journal of Clinical Nutrition*, 84(2), 289-298.
- Shivaramakrishna, H., Wantamutte, A., Sangolli, H., & Mallapur, M. (2010). Risk factors of coronary heart disease among bank employees of Belgaum city. A Cross-sectional study. Al Ameen Journal of Medical Sciences, 3(2), 152-159.
- Singh, S., Shankar, R., & Singh, G. P. (2017). Prevalence and associated risk factors of hypertension: A cross-sectional study in urban Varanasi. *International Journal of Hypertension*, 2017, 1-10.
- Sovova, E., Nakladalová, M., Kaletova, M., Sovova, M., Radova, L., & Kribska, M. (2014).
 Which health professionals are most at risk for cardiovascular disease? Or do not be a manager. *International Journal of Occupational Medicine and Environmental Health*, 27(1), 71-77.
- Sun, Z. (2015). Aging, arterial stiffness, and hypertension. Hypertension, 65(2), 252-256.
- Szapary, P. O., Bloedon, L. T., & Foster, G. D. (2003). Physical activity and its effects on lipids. *Current Cardiology Reports*, 5(6), 488-493.
- Ulasi, I. I., Ijoma, C. K., & Onodugo, O. D. (2010). A community-based study of hypertension and cardio-metabolic syndrome in semi-urban and rural communities in Nigeria. BMC Health Services Research, 10(1), 1-6.
- Van de Vijver, S., Akinyi, H., Oti, S., Olajide, A., Agyemang, C., Aboderin, I., & Kyobutungi,
 C. (2014). Status report on hypertension in Africa-Consultative review for the 6th
 Session of the African Union Conference of Ministers of Health on non-communicable
 diseases. *Pan African Medical Journal*, 16(1), 38-55.
- Vorster, H. (2002). The emergence of cardiovascular disease during urbanisation of Africans. *Public Health Nutrition*, *5*(1a), 239-243.

- Walker, R. W., McLarty, D. G., Kitange, H. M., Whiting, D., Masuki, G., Mtasiwa, D. M., Machibya, H., Unwin, N., & Alberti, K. M. (2000). Stroke mortality in urban and rural Tanzania. *The Lancet*, 355(9216), 1684-1687.
- Wamala, J. F., Karyabakabo, Z., Ndungutse, D., & Guwatudde, D. (2009). Prevalence factors associated with hypertension in Rukungiri district, Uganda-a community-based study. *African Health Sciences*, 9(3), 1-12.
- Werneck, A. O., Oyeyemi, A. L., Gerage, A. M., Cyrino, E. S., Szwarcwald, C. L., Sardinha, L. B., & Silva, D. R. (2018). Does leisure-time physical activity attenuate or eliminate the positive association between obesity and high blood pressure? *The Journal of Clinical Hypertension*, 20(5), 959-966.
- WHO. (2005). The WHO STEPwise Approach to Chronic Disease Risk Factor Surveillance https://apps.who.int/iris/handle/10665/43376.
- WHO. (2009). 2008-2013 Action plan for the global strategy for the prevention and control of non-communicable diseases: Prevent and control cardiovascular diseases, cancers, chronic respiratory diseases and diabetes. World Health Organization. https://www. who.int/nmh/publications/ncd_action_plan_en.pdf.
- WHO. (2010). Healthy workplaces: A model for action-for employers, workers, policy-makers and practitioners. World Health Organization. https://apps.who.int/iris/handle/10665/ 44307.
- WHO. (2013). A global brief on hypertension: Silent killer, global public health crisis: World Health Day 2013. https://apps.who.int/iris/handle/10665/79059.
- WHO. (2014). Prevalence of Noncommunicable Disease Risk Factors in the Republic of Moldova Regional Office for Europe. https://www.euro.who.int/__data/assets/pdf_ file/0009/252774/PREVALENCE-OF-NONCOMMUNICABLE-DISEASE-RISK-FACTORS-IN-THE-REPUBLIC-OF-MOLDOVA-STEPS-2013.pdf
- WHO. (2016). *Global database on body mass index*. 2006. https://www.euro.who.int/en/ health-topics/disease prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi.

- WHO. (2018). Non-communicable diseases country profiles 2018. World Health Organization. https://apps.who.int/iris/handle/10665/274512.
- Wijayathunge, L., & Hettiaratchi, U. (2017). Assessment of knowledge and risk factors of hypertension among school teachers in a selected district in North Central Province of Sri Lanka. *International Journal of Scientific and Research Publications*, 7(11), 1-5.
- Williams, E. A., Keenan, K. E., Ansong, D., Simpson, L. M., Boakye, I., Boaheng, J. M., Awuah, D., Nkyi, C. A., Nyannor, I., & Arhin, B. (2013). The burden and correlates of hypertension in rural Ghana: A cross-sectional study. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*, 7(3), 123-128.
- Worrall, L., Colenbrander, S., Palmer, I., Makene, F., Mushi, D., Mwijage, J., Martine, M., & Godfrey, N. (2017). Better urban growth in Tanzania: Preliminary exploration of the opportunities and challenges. In: Coalition for Urban Transitions, London and Washington, DC. http://newclimateeconomy. .net/content/cities-working-papers.
- Yanagiya, S., Nakamura, K., Ukawa, S., Tsutsumi, A., Atsumi, T., & Tamakoshi, A. (2020).
 Household income and the risk of incident hypertension in employees at multiple workplaces in Japan: J-HOPE. *Hypertension Research*, 43(12), 1445-1453.
- Yu, Z., Nissinen, A., Vartiainen, E., Song, G., Guo, Z., Zheng, G., Tuomilehto, J., & Tian, H. (2000). Associations between socio-economic status and cardiovascular risk factors in an urban population in China. *Bulletin of the World Health Organization*, 78, 1296-1305.
- Zhou, B., Bentham, J., Di Cesare, M., Bixby, H., Danaei, G., Cowan, M. J., Paciorek, C. J., Singh, G., Hajifathalian, K., & Bennett, J. E. (2017). Worldwide trends in blood pressure from 1975 to 2015: A pooled analysis of 1479 population-based measurement studies with 19. 1 million participants. *The Lancet*, 389(10064), 37-55.

APPENDICES

Appendix 1: Structured questionnaire for the assessment of risk factors for hypertension and assessment of work-related stress

(a) Structured questionnaire for the assessment of risk factors for hypertension

Questionnaire ID.....

INSTRUCTIONS: Please would you put $(\sqrt{)}$ in an appropriate box according to the answer selected.

	STEP I. DEMOGRAPHIC INFORMATION			
Den	nographic information			
	Question	Response		
1.	Sex	Male []		
		Female []		
2.	How old are you?	years old		
3.	What is the highest level of education you have completed?	 [] Up to primary school [] Secondary level [] College/University completed [] Post graduate degree 		
4.	Job profession			
5.	What is your marital status?	 [] Married/Cohabiting [] Single [] Separated/Divorced [] Widowed 		
6.	Which of the following best describes your main work status over the past 12 months?	[] Government employee [] Non-government employee		

7.	How long have you been working	[] Less than a year	
	with the current institution?	[] 1-3 years	
		[] 4-6 years	
		[] 7 years and above	
8.	How many people older than 18 years, including yourself, live in your household?	Number	
9.	What is your estimated monthly	[]≤250 000	
	income? (In Tanzanian shillings)	[] > 250 000 to < 500 000	
		[] 500 000 to < 750 000	
		[] 750 000 to < 1 000 000	
		[] > 1 000 000	
STE	CP I. BEHAVIORAL MEASURE	MENTS	
I am like Tob	going to ask you some questions al smoking, drinking alcohol, eating fracco use	bout various health behaviours. This i ruits and vegetables and physical activ	ncludes things vity
10.	Do you currently smoke any	[]Yes	If no go to
	tobacco products, such as cigarettes, cigars or pipes daily?	[] No	question number 13
11.	How old were you when you first started smoking daily?	years old	
12.	On average, how many of the	Number []	
	following do you smoke each day?	[] Manufactured cigarette	
		[] Hand-rolled cigarettes	
		[] Pipes full of tobacco	
		[] Cigars (cheroots)	

13.	In the past, did you ever smoke tobacco daily?	[] Yes [] No	If no go to question number 15
14.	How long ago did you stop tobacco smoking daily?	[] Years ago[] Months ago[] Weeks ago	
15.	Do you currently use smokeless tobacco products daily?	[] Yes [] No	
16.	In the past, did you ever smoke daily?	[] Yes [] No	
17.	During the past 30 days, on how many days did someone smoke when you were present either at home or around your workplace?	Number of daysHome[Working place[
Alco	ohol consumption	[
	Question	Response	
18.	Have you ever consumed an alcoholic drink such as beer, wine, spirits and homebrewed wine?	[] Yes [] No	If no go to question number 23
19.	In the past 12 months have you consumed an alcoholic drink?	[] Yes [] No	
20.	Have you consumed an alcoholic drink within the past 30 days?	[] Yes [] No	
21.	During the past 30 days, when you drank alcohol, on average, how many standard alcoholic	Number []	

	drinks did you have during one drinking occasion?		
22.	During the past 30 days, when you consumed an alcoholic drink, how often was it with meals? Please do not count snacks.	 Usually with meals Sometimes with meal Rarely with meals Never with meals 	

Diet

The next questions ask about the fruits and vegetables that you usually eat. I have a nutrition card here that shows you some examples of local fruits and vegetables. Each picture represents the size of a serving.

	Question	Response	
23.	In a typical week, on how many days do you eat fruit and	Number of days	
	vegetables?	Vegetables []	
24.	How many servings of fruits and vegetables do you eat on one of those days?	Number of servings Fruits []	
		Vegetables []	
25.	What type of oil or fat is most often used for meal preparation in your household?	[] Safi [] Korie	
		[] Sunola	
		[] Sunflower	
		[] Olive oil	
		[] Margarine	
		Others (specify)	

26.	On average, how many meals	Number of meals	
	per week do you eat that were not prepared at home?	Breakfast []	
		Lunch []	
		Dinner []	
Diet	ary salt		
	Question	Response	
27.	How often is salt, salty seasoning and sauces added in	[] Always	
	cooking or during food preparation at home?	[] Often	
	Propulsion of notice .	[] Sometimes	
		[] Rarely	
		[] Never	
28.	How often do you add salt or salty sauce such as ketchup or	[] Always	
soya	soya sauce to your food before	[] Often	
	you eat it or while eating?	[] Sometimes	
		[] Rarely	
		[] Never	
29.	Do you always use table salt	[] Yes	
	while eating	[] No	
30.	What kind of salt do you use for cooking or meal preparation in	[] Iodized	
	your household?	[] Non-iodized	
		[] Don't know	
31.	How often do you eat processed food (the foods that have been	[] Always	

32.	altered from their natural state, such as packaged snacks, canned food, fast-food (e.g. pickles, marinates, sheep's cheese, salami, sausages, ham and other meat products, salted nuts/biscuits, cakes, Crips)) How much salt is recommended by WHO per day	 [] Often [] Sometimes [] Rarely [] Never [] Too much [] 10g [] 5g 	
		[] 15g	
		[] Don't know	
33.	How much salt or salty sauce do you think you consume?	 Right amount Too little Don't know 	
34.	Do you think reducing the amount of salt you add to the food is important?	[] Yes [] No [] I don't know	
35.	Do you think that too much salt or salty sauce in your diet could Cause a health problem?	[] Yes [] No [] I don't know	
36.	If yes in above what sort of health problem do you think can be caused by a high salt diet? Put a tick to all that apply	 High blood pressure Osteoporosis Stomach cancer Kidney stones Heart attack/ failure Stroke 	

		 [] Asthma [] Don't know [] Other (specify) 	
37.	Do you do have regular strategies to control your salt intake?	[] Yes [] No If yes indicate	

Physical activities at Work

Vigorous-intensity activity are the activities that causes large increases of heart rate like lifting heavy loads, manual construction work, while moderate-intensity activity causes small increases of heart rate such as brisk walking, carrying light loads, and light activities are the activities that that does not increase the heart beat or heart rate like sitting in the office.

	Question	Response	
38.	Does your work involve either of the following activity at least 10 minutes continuously?	 [] Vigorous-intensity activity [] Moderate-intensity activity [] Light activity 	
39.	How many days do you do the activity selected above as part of your work in a week?	Number of days []	
40.	How much time do you spend doing the activities at work on a typical day?	Number of Hours []: Minutes []	
Travel to and from places			

The usual way to travel to and from places include travel for example to work, for shopping, to market and the place of worship.

Question	Response	

41.	Which means of transport do you use continuously from home to working place? In a typical week, how many days and time do you spend	 [] Bicycle (pedal cycle) [] Walking [] Car [] Motorcycle Number of 	
	walking or bicycle to get to and from places?	Day [] Hours [] Minutes []	
Rec	reational activities		
This trans	asks you about sports, fitness and r sport activities that you have already	ecreational activities (leisure) excludi y mentioned	ing work and
	Question	Response	
43.	Do you do vigorous-intensity sports or recreational activities that cause large increases in breathing or heart rate such as running or football, for at least 10 minutes continuously?	[] Yes [] No	
44.	In a typical week, on how many days do you do vigorous- intensity sports, fitness or recreational activities?	Number of days []	
45.	How much time do you spend doing vigorous-intensity sports, fitness or	Number of Hours []: Minutes []	
	Recreational activities on a typical day?		
46.	Do you do moderate-intensity sports or recreational activities that cause a	[] Yes [] No	

	a small increase in breathing or heart rate, such as brisk walking, cycling, Swimming, volleyball for at least 10 minutes continuously?		
47.	In a typical week, on how many days do you do moderate- intensity sports, fitness or recreational activities?	Number of days []	
48.	How much time do you spend doing Moderate-intensity sports, fitness or recreational (leisure) activities on a typical day?	Number of Hours []: Minutes []	
49.	How much time do you usually spend sitting or reclining for example watching Television, listening to the music, reading books or newspaper without including the time spent sleeping.	Number of Hours []: Minutes []	
Hist	ory of Hypertension, Diabetes and	d Blood cholesterol	
	Question	Response	
50.	Have you ever had your blood pressure blood cholesterol and blood glucose measured by a doctor or other health worker?	HypertensionDiabetesBloodcholesterolYes [][]No [][]	
51.	Have you ever been told by a doctor or other health worker that you have hypertension or diabetes?	Hypertension Diabetes Blood cholesterol Yes [] [] [] No [] [] []	
52.	Are you currently taking any drugs (medication) in the past	Hypertension Diabetes Blood cholesterol	

	two weeks for hypertension, blood cholesterol or diabetes?	Yes [] [] []				
		No[] [] []				
53.	Are you currently taking any herbal or traditional remedy for	Hypertension Diabetes Blood cholesterol				
	hypertension or diabetes?	Yes [] [] []				
		No[] [] []				
54.	In your family do you have any	Diabetes []				
	diabetes or hypertension or both	Hypertension []				
		Both of them []				
STE	CP II: PHYSICAL MEASUREME	INT				
Hei	ght and weight					
	Question	Response				
55.	Height	In centimeter (cm)				
56.	Weight	In kilogram (kg)				
Wai	st circumference					
57.	Waist circumference	In Centimeter (cm)				
Bloc	Blood pressure measurement					
58.	Reading 1	Systolic (mmHg)				
		Diastolic (mmHg)				
	Reading 2	Systolic (mmHg)				
		Diastolic (mmHg)				
	Reading 3	Systolic (mmHg)				

		Diastolic (mmHg)		
Bloc	od measurement			
Bloc	od glucose			
59.	Fasting blood glucose	mmol/l		
Blood cholesterol				
60.	Total cholesterol	mmol/l		
	Low-Density Lipoprotein(LDL- C)	mmol/l		
	High-Density Lipoprotein(HDL- C)	mmol/l		
	Triglyceride	mmol/l		

(b) Assessment of work-related stress

ASS	ESSMENT OF WORK RELATED STRESS					
61.	I have unachievable deadlines	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
62.	I have to work very intensively	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
63.	I have to neglect some tasks because I have too much	Never	Seldom	Sometimes	Often	Always
	to do	[]	[]	[]	[]	[]
64.	I am unable to take sufficient breaks	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
65.	I am pressured to work long hours	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
66.	I have to work very fast	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
67.	I have unrealistic time pressures	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
68.	I can decide when to take a break	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
69.	I have a say in my work speed	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
70.	I have a choice in deciding how I do my work	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
71.	My working time can be flexible	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
72.	If work gets difficult, my colleagues will help me	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
73.	I am given supportive feedback on the work I do	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
74.	I get the help and support I need from colleagues	Never	Seldom	Sometimes	Often	Always
		[]	[]	[]	[]	[]
75.	I receive the respect at work I deserve from my	Never	Seldom	Sometimes	Often	Always
	colleagues	[]	[]	[]	[]	[]

76.	I can talk to my line manager about something that	Never Seldom Sometimes Often Always
	has upset or annoyed me about work	
77.	My colleagues are willing to listen to my work-	Never Seldom Sometimes Often Always
	related problems	
78.	I am supported through emotionally demanding work	Never Seldom Sometimes Often Always
79.	My line manager encourages me at work	Never Seldom Sometimes Often Always
80.	Is the salary enough to meet the daily basic need?	[]Yes []No
81.	Do you experience an unplanned shift	[]Yes []No
82.	Do you experience long working hours	[] Yes [] No
83.	Poor working condition e.g. excessive noise,	[] Yes [] No
	temperature extremes, dust etc	
84.	Long-distance from home to work	[]Yes []No
85.	Do you feel stressed at work	[]Yes []No
86.	What is your level of stress	Very low [] Low [] Moderate [] High [] Very high
		[]

Frequency						
Food item	Never	Rarely	Everyday	1-2	3-4	1-3
		(≤ than		per	per	per
		once a		week	week	month
		month)				
Cereals and cereal products						
Maize on cob						
Mix of beans and other cereals						
Porridge						
Rice						
Breakfast cereals						
Stiff porridge						
Spaghetti						
Banana, Roots and tubers		<u>.</u>	-			
Cooked Banana						
Fried Banana						
Cassava						
Sweet potato						
Arrowroot						
Potatoes						
French fries						
Banana stew						
Crips						
Meat, fish and animal product	t	<u>.</u>	-			
Beef						
Mutton						
Organ meat						
Pork						
Chicken						
Sausage						
Egg						
Fish						
Sardines						
Milk						
Yoghurt						
Legumes and nuts			1		1	
Beans						
Green peas						
Bambara nuts						
Groundnuts						
Cashew nuts						

Appendix 2: Assessment of dietary practices

Fruits					
Cucumber					
Banana					
Orange					
Watermelon					
Avocado					
Pineapple					
Baobao					
Mango					
Vegetables					
Cabbage					
Amaranth leaves					
Spinach					
Cowpea leaves					
Pumpkin leaves					
African eggplant					
African nightshade					
Carrot					
Tomato					
Fats and oil					
Vegetable fat					
Mayonnaise					
Lard					
Peanut butter					
Other foods					
Pizza					
Burger					
Water					
Carbonated soft drink					
Alcoholic drink					
Coffee					
Black tea					
Milk tea					

Appendix 3: Showcard

(Diet) Serving Sizes of Fruits and Vegetables

VEGETABLES ARE	1 SERVING	EXAMPLES
CONSIDERED TO BE:	(STANDARD)	
Raw green leafy vegetables	1 cup	Spinach, salad, etc.
Other vegetables, cooked or chopped raw	¹ /2 Cup	Tomatoes, carrots, pumpkin, cabbage, fresh beans, onion, Etc
FRUIT		
Cucumber (Matango)	1 medium-size piece	
Banana (Ndizi)	1 medium-size piece or 2 small bananas(ndizi kisukari)	

Orange (Chungwa)	1 medium-size piece	
Mango (Embe)	1 medium-size piece	
Watermelon (Tikiti maji)	¹ ⁄4 Kipande cha tikiti maji (134g)	Weitermeton
Avocado (Parachichi)	1 small size (parachichi dogo) (110 g)	
Pawpaw (Papai)	¹ /4 piece (¹ /4 kipande) (120g)	
Pineapple (Nanasi)	¹ ⁄4 piece	
Fruit juice	1/2 cup	Juice from fruit.
Appendix 4: Informed consent form

THE NELSON MANDELA AFRICAN INSTITUTION OF SCIENCE AND TECHNOLOGY (NM-AIST)



CONSENT FORM

TITLE: DETERMINATION OF FACTORS ASSOCIATED WITH HYPERTENSION AMONG EMPLOYEES IN ARUSHA CITY COUNCIL, TANZANIA.

This Informed Consent Form has two parts:

- Part I: Information sheet
- Part II: Certificate of Consent (for signatures if you agree to take part)

PART I: INFORMATION SHEET

Introduction: I am Dalahile Zubery, a master student from the Nelson Mandela African Institution of Science and Technology (NM-AIST). I am currently involved in conducting research on the Determination of Factors Associated with Hypertension among Employees in Arusha City Council. Please listen carefully and ask any questions you may have before agreeing to take part in the study.

What the study is about: The purpose of this study is to assess the factors associated with hypertension among employees in Arusha City Council, Tanzania.

Purpose of the research: Hypertension is affecting the majority of the people in the country and it has become among the ten diseases that contribute to death in the country. There is a need for identifying the risk factors associated with hypertension and make strategies to mitigate the risk factors.

Participant Selection: You are being invited to take part in this research because you are the employee falling in the category group that demonstrated to be at risk of the disease. If you become affected with the disease it may result in loss of manpower due to repeated absenteeism

at workplaces which can result in early retirement but there is also loss of income through providing care for the management of the disease.

What we will ask you to do: If you agree to be in this study, I will conduct a survey with you. The survey will include questions about your lifestyle behaviours, dietary habit, and measurement on body weight, height and waist circumference but also a collection of (5ml) of a blood sample for analysis of blood glucose, total blood cholesterol, high-density lipoprotein, low-density lipoprotein, and triglyceride the survey will take about 90 minutes to complete.

Risks: There is the risk that you may find some of the questions about your household to be sensitive. You may refuse to answer any particular question and may stop the interview at any time.

Benefits: The immediate benefits to you for taking part in this survey is that you will be able to know your health status and get free medical advice based on the screening result but the longer-term benefit will aim at developing workplace intervention strategies to mitigate the risk factors for hypertension and related NCDs.

Compensation: There is no compensation for taking part in this study.

Confidentiality: Your answers will be confidential. The records of this study will be kept private. In any sort of report, we make public we will not include any information that will make it possible to identify you. Research records will be kept in a locked file; only the researchers will have access to the records.

Taking part is voluntary: Taking part in this study is completely voluntary. You may skip any questions that you do not want to answer. If you decide not to take part or to skip some of the questions, it will not affect your current or future relationship with Nelson Mandela African Institute of Science and Technology, or the District Council. If you decide to take part, you are free to withdraw at any time.

PART II: CERTIFICATE OF CONSENT

I have read the foregoing information or it has been read to me and has understood. My questions have been answered to my satisfaction. I agree to participate in this study.

Name of participant _____

Signature of participant _____ Date _____

I confirm that the participant was allowed to ask questions about the study, and all the questions asked by the participant have been answered correctly and to the best of my ability. I confirm that the individual has not been coerced into giving consent, and the consent has been given freely and voluntarily.

Name of researcher _____

Signature of researcher _____ Date _____

Thank you very much for your participation in this study.

Appendix 5: KNCHREC ethical clearance



Poster Presentation



FACTORS ASSOCIATED WITH HYPERTENSION AMONG EMPLOYEES IN ARUSHA CITY, TANZANIA

Dalahile Zubery ^{1*}Judith Kimiywe ² Haikael D Martin¹

Abstract

Healthcare workers, teachers and bankers are occupation groups exposed to hypertension due to the nature of their work. A descriptive cross-sectional study was conducted to identify the predicting factors for hypertension among teachers, bankers and healthcare workers in Arusha city council. The prevalence of hypertension among working adults was 23.8%. Age, alcohol consumption, low salary and high Low-Density Lipoprotein cholesterol (LDL-C), were significantly associated with hypertension.

Background

Hypertension is the major risk factor for CVDs, like heart attack, stroke and heart failure. In a review of hypertension prevalence in developing countries, a higher hypertension prevalence was reported in Tanzania than in many other developing countries. Despite that, there is limited documentation on the predictors of hypertension among working adults in Tanzania. Therefore, the current study aimed to identify the factors associated with hypertension among healthcare workers, teachers and bankers in Arusha City Council.

Results

The overall hypertension prevalence among working adults was 23.8%. Age (AOR=34.98, 95% CI: 1.30-94.03), alcohol consumption (AOR=6.55, 95% CI: 1.22-35.28), low salary (AOR=6.44, 95% CI: 1.12-37.18) and high Low-Density Lipoprotein cholesterol (LDL-C) (AOR=5.93, 95% CI: 1.24-28.45), were significantly factors associated with hypertension.

Conclusion

The current study found high prevalence of hypertension, overweight and obesity among working adults and the higher prevalence bankers was among compared with other occupations. The finding shows an emerging problem in the working population, especially among bankers compared to other occupations. Hence this findings inform the policymakers, education, financial and health sectors to design workplace wellness program specifically on healthy lifestyle to mitigate this factors.

Acknowledgement

African Center for Research, Agriculture advancement Teaching Excellence and Sustainability (CREATES) at Nelson Mandela-African Institution of Science and Technology for financial support

Author's affiliations

¹School of Life Sciences and Bioengineering (LiSBE), The Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania.

²Department of Foods, Nutrition and Dietetics, Kenyatta University, Nairobi, Kenya.