

2024-11-25

Enhancing the optoelectronic properties of blended triphenylamine-betalain based dyes through tailoring the anchoring unit: a theoretical investigation

Tsere, Melkizedeck

Taylor & Francis

<https://doi.org/10.1080/00268976.2024.2429752>

Provided with love from The Nelson Mandela African Institution of Science and Technology

Enhancing the optoelectronic properties of blended triphenylamine-betalain based dyes through tailoring the anchoring unit: a theoretical investigation

Melkizedeck Hiiti Tsere, Rene Costa, Geradius Deogratias, Ohoud S Al-Qurashi, Nuha Wazzan, Tatiana Pogrebnaya, Alexander Pogrebnoi, Revocatus Machunda

To down the complete text, click that link.

DOI: <https://doi.org/10.1080/00268976.2024.2429752>

Abstract

A series triphenyl-betalain organic dyes featuring carboxylic acid and nitro anchoring groups $\text{CH}=\text{C}(\text{X})\text{COOH}$ for the A1-X dyes and $-\text{CH}=\text{C}(\text{X})\text{NO}_2$ for the A2-X dyes, respectively, where $\text{X}=\text{CN}$, CH_3 , CCl_3 and CF_3 was evaluated for dye sensitised solar cells application. The geometrical structures, molecular orbitals and energies, light absorption patterns, free energies of electron injection and dye regeneration and binding to the semiconductor have been explored using DFT/TD-DFT methods. The nitro-based anchoring group resulted in pronounced red-shift in absorption spectra between 111 and 317 nm compared to carboxylic acid-based dyes. Attachment of the dyes to the semiconductor was modelled via binding to $(\text{TiO}_2)_6\text{H}_3$ cluster; A2-X dyes exhibited more stable Dye@TiO₂ complexes with binding energies (BEs) ranging between -4.08 and -2.88 eV compared to A1-X dyes with BEs range of -1.11 to -0.05 eV. The results evince that the dyes with $\text{CH}=\text{C}(\text{X})\text{NO}_2$ anchoring groups could be promising materials for light harvesting application.

Keywords; Anchoring group, DFT, betalain, triphenyl-betalain, sensitiser